

New Mn II energy levels from the STIS-HST spectrum of the HgMn star HD 175640[★]

F. Castelli¹, R. L. Kurucz², and C. R. Cowley³

¹ Istituto Nazionale di Astrofisica, Osservatorio Astronomico di Trieste, Via Tiepolo 11, I-34143 Trieste, Italy

e-mail: castelli@oats.inaf.it

² Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA

e-mail: rkurucz@cfa.harvard.edu

³ Department of Astronomy, University of Michigan, Ann Arbor, MI 48109-1042, USA

e-mail: cowley@umich.edu

ABSTRACT

Aims. The NIST database lists several Mn II lines that were observed in the laboratory but not classified. They cannot be used in spectrum synthesis because their atomic line data are unknown. These lines are concentrated in the 2380-2700 Å interval. We aimed to assign energy levels and log *gf* values to these lines.

Methods. Semiempirical line data for Mn II computed by Kurucz were used to synthesize the ultraviolet spectrum of the slow-rotating, HgMn star HD 175640. The spectrum was compared with the high-resolution spectrum observed with the HST-STIS equipment. A UVES spectrum covering the 3050-10000 Å region was also examined.

Results. We determined a total of 73 new energy levels, 58 from the STIS spectrum of HD 175640 and another 15 from the UVES spectrum. The new energy levels give rise to numerous new computed lines. We have identified more than 50% of the unclassified lines listed in the NIST database and have changed the assignment of another 24 lines. An abundance analysis of the star HD 175640, based on the comparison of observed and computed ultraviolet spectra in the 1250-3040 Å interval, is the by-product of this study on Mn II.

Key words. line:identification-atomic data-stars:atmospheres-stars:chemically peculiar- stars:individual: HD 175640

1. Introduction

The availability of a reliable and complete set of atomic data is an essential requirement for research in stellar physics, in particular for computing stellar atmospheres and stellar spectra to be compared with observations at various resolutions.

The high-resolution, high signal-to-noise stellar spectra observed by the modern telescopes and spectrographs have proved to be a useful tool for studies aimed at extending the knowledge of atomic and molecular data. An example is the analysis of the UVES spectrum of the peculiar star HR 6000 which has permitted us to fix about 120 Fe II energy levels not observed before in the laboratory, but predicted with approximate energy values by the theory (Castelli et al. 2008, Castelli et al. 2009, Castelli & Kurucz 2010). Another example of the use of stellar spectra to derive atomic data is the paper by Peterson & Kurucz (2015) who obtained 66 new energy levels for Fe I from the analysis of HST-STIS spectra of a group of 13 metal-poor stars. In the above works, for each new determined energy level, several new transitions with wavelengths and oscillator strengths are associated, so that the line lists used for computing spectra are considerably augmented.

In this paper we extend to Mn II the above described studies. History has shown that the refinement and extension of our atomic database has lead to important astronomical discover-

ies. In the case of the Mn II spectrum, the subject of the present paper, the extended analysis by Iglesias & Velasco (1964) enabled many new line identifications in the class of stars now known as mercury-manganese (HgMn) stars, occupying the spectral region ~ A0/B9-B6 (10500-16000K). This in turn enabled Bidelman (1962) and Dworetzky (1969) to show that these stars were enormously enriched in mercury and platinum. In the present study, we have been able to obtain abundances for ten elements that were unavailable from the ground spectra (B, N, Al, Cl, V, Zn, Ge, As, Ag, and Cd). The ability to distinguish their spectral lines from those of Mn II was essential.

To study the Mn II spectrum, we adopted the same procedure described by Castelli & Kurucz (2010). We analysed HST-STIS spectra from 1250-3045 Å and a UVES spectrum from 3050 to 10000 Å of the HgMn star HD 175640. It is a slowly rotating star with an abundance in manganese on the order of 2.5 dex over the solar value. The UVES spectrum, previously analyzed to derive stellar parameters and abundances by Castelli & Hubrig (2004), was further examined for this paper to derive new Mn II energy levels.

To fix new energy levels we adopted the semiempirical Mn II line data computed by Kurucz. We determined a total of 73 new energy levels, 58 from the HST-STIS spectrum and 15 from the UVES spectrum. Updated line lists for most elements were adopted in order to synthesize the spectra. The most recent work on Mn II is that of Kramida & Sansonetti (2013) who critically analyzed all the atomic data available in the literature for this ion. They revised the Sugar & Corliss (1985) energy levels and provided a list of Mn II spectral lines and transition probabili-

Send offprint requests to: F. Castelli

[★] Tables A.1 and A.2 are only available in electronic form at CDS via anonymous ftp to cdsarc.u-strasbg.fr (130.79.128.5) or via <http://cdsweb.u-strasbg.fr/cgi-bin/qcat?J/A+A/>

ties taken mostly from experimental sources. The Mn II atomic data of the NIST database (Kramida et al. 2014) are the result of the Kramida & Sansonetti (2013) work. In the database there are numerous unclassified lines of Mn II. They are concentrated in the 2380–2700 Å interval which was observed for HD 175640 at about 120000 resolution. The new Mn II energy levels have permitted us to identify more than 50% of the lines unclassified in the NIST database.

2. Observations and data reduction

HD 175640 (HR 7143) is one of the targets included in the “Hot Stars” program (GO-13346), which is part of the “Advanced Spectral Library (ASTRAL)” Project (Ayres 2014). The star was observed in Cycle 21 of the Space Telescope Imaging Spectrograph (STIS). The spectrum covers the range 1150–3045 Å. The nominal resolving power R ranges from about 30000 to 120000; the S/N ratio is larger than 100. We used the final spectrum that resulted from the calibration and the merging of the individual spectra observed in the different wavelength intervals, as performed by the ASTRAL Science Team (Carpenter et al. 2014).

For this work we analyzed the whole region from 1250 to 3045 Å. We used the IRAF tool “continuum” to normalize the observed spectrum to the continuum level. When we compared observed and computed spectra we tentatively fixed different resolving powers for the different spectral intervals roughly corresponding to the different resolutions of the different observed regions (Ayres 2010). In particular, we adopted R equal to 50000, 40000, 25000, 30000, and 120000 for the ranges 1250–1340 Å, 1340–1690 Å, 1690–2200 Å, 2200–2332 Å, and 2332–3045 Å, respectively. The observed spectrum was shifted to the Ritz wavelengths, as derived from the energy levels, using mostly Fe II lines as reference lines, owing to their accurate wavelengths derived from the Nave & Johansson (2013) energy levels. The shift to superimpose the observed spectrum to the synthetic spectrum is $+25 \text{ km s}^{-1}$ in the 1250–2000 Å region. After conversion of the wavelength scale of the 2000–3040 Å interval from vacuum to air, the shift is $+24.5 \text{ km s}^{-1}$. We estimate an uncertainty in the velocity shift on the order of 0.5 km sec^{-1} , corresponding to an uncertainty in the wavelength scale of 0.2 mÅ , 0.35 mÅ , and 0.5 mÅ at 1250 Å, 2000 Å, and 3000 Å, respectively.

3. The star HD 175640

The stellar parameters and the abundances of HD 175640 were determined by Castelli & Hubrig (2004) in a previous analysis of the optical spectrum of the star. An ATLAS12 (Kurucz 2005) model with parameters $T_{\text{eff}}=12000 \text{ K}$, $\log g=3.95$, and microturbulent velocity $\xi=0 \text{ km sec}^{-1}$ was computed for the individual stellar abundances. They were obtained from the equivalent widths of selected lines and are listed in the third column of Table 1. The SYNTH code (Kurucz 2005) was used to compute a synthetic spectrum for the 3050 – 10000 Å region which was compared with the observed spectrum. The ATLAS12 model and the given set of abundances were adopted. A rotational velocity of $v \sin i = 2.5 \text{ km sec}^{-1}$ was derived from the comparison of the observed and computed spectra. The adopted zero microturbulent velocity was based both on the Adelman (1994) conclusions that most HgMn stars have little or no microturbulent velocity and on the consistent abundances we obtained from the equivalent widths of 55 weak and strong Fe II lines measured in the optical region (Castelli & Hubrig 2004).

In this paper we compared the observed HST-STIS spectrum with a synthetic spectrum computed for the 1250–3045 Å interval. We used the ATLAS12 model atmosphere adopted in Castelli & Hubrig (2004) and the abundances listed in the second column of Table 1. We started with the abundances adopted to compute the optical spectrum and then we modified some of them in order to better fit the ultraviolet spectrum. The abundances of B, N, Al, Cl, V, Zn, Ge, As, Ag, and Cd were derived only in this paper. For those elements that do not show lines in the HST-STIS spectrum we adopted the abundances derived from the optical region if they did not produce inconsistent results, as is the case of Pt. For the ions Cr I, Pt I, Ga III, Y III, and Yb III the ionization equilibria are not satisfied. We did not use their corresponding abundance to compute the synthetic spectrum, but that of the ions in the other ionization stages. In addition, the abundance derived from Mn I and Mn II gives incorrect profiles for a large number of Mn III lines, that are either not observed or are computed as too strong (Figure 1). These lines are mostly due to transitions having either an even level with energy larger than 172000 cm^{-1} or an odd level with energy larger than 130000 cm^{-1} . We compared the Kurucz $\log gf$ values that we used with those from Uylings & Raassen (1997) for the lines in common. Because the differences are not larger than 0.2 dex (while fitting the observed lines requires lowering the $\log gf$ values by 1.0 up to 2.0 dex or more), we could argue that some physical mechanism like the vertical abundance stratification weakens the high-excitation lines of Mn III in HD 175640. On the other hand, the low-excitation lines, in particular those of multiplets 10, 11, 12, 13, 14, 15, 16, 17, and 18 (Moore 1950), which lie in the 1900–2400 Å region, are reasonably well predicted by the adopted abundance.

To compute the synthetic spectrum we used the Kurucz database which is formed by the Kurucz (1988; 1993) line lists and by new and revised data (Kurucz 2011). The Kurucz line lists include the hyperfine structure for several elements. In particular, for Mn II, the magnetic dipole and electric quadrupole coupling constants A and B measured by Holt et al. (1999) for 59 levels were used to compute the hyperfine structure of all the lines with coupling constants available for both levels. The constants and the hyperfine lines are listed on Kurucz’s website¹. The two files quoted in the footnote differ in that experimental $\log gf$ values, when available, are used in the second file instead of the Kurucz computed values which are stored in the first file.

For this work we added lines of Ga II, Ga III, Rh II, Pd II, Ag II, Y III, Zr III, Yb III, Pt II, Pt III, Au II, Au III, Hg II, Hg III. The presence of these elements in the STIS spectrum of HD 175640 was inferred from the previous analysis of the optical region (Castelli & Hubrig 2004). We also used predicted lines and intensities for the STIS region using the “predicted stellar” option of the VALD database (Ryabchikova et al. 1997) with stellar parameters for HD 175640 from Castelli & Hubrig (2004). Additionally, we used results from wavelength coincidence statistic (WCS, Cowley & Hensberge 1981), using measurements of STIS wavelengths for $\lambda > 2000 \text{ Å}$. These wavelengths are available online². More information on the heavy elements identified in HD 175640 is given in the Appendix.

In addition to the large overabundance of Mn ([+2.4]), Ti ([+1.4]), Cr ([+1.0]), and the mild underabundance of Fe ([−0.3]), mild overabundances of B, Na, and P, a general underabundance of the light elements (He, C, N, Mg, Al, Si, S,

¹ <http://kurucz.harvard.edu/atoms/2501/hyper250155.pos> or [hyper250155.all](http://kurucz.harvard.edu/atoms/2501/hyper250155.all)

² <http://dept.astro.lsa.umich.edu/~cowley/HR7143air.html>




Fig. 1. Example of computed spectrum (red line) displaying Mn II lines not observed in the stellar spectrum (black line). The lines are those at 1605.126 Å, 1605.545 Å, and 1605.690 Å. Their upper level is an even level with energy 179152.090 cm⁻¹, 172682.640 cm⁻¹, and 172451.230 cm⁻¹, respectively. The labels for the lines include the last three digits of the wavelength, the code (atomic number) for the element and ion, the lower energy level in cm⁻¹, and the residual intensity at line center in per mil.

Cl), and general overabundance of several heavy elements (Cu, Ga, As, Br, Sr, Y, Zr, Rh, Pd, Ag, Cd, Xe, Yb, Au, and Hg) were measured. This peculiar chemical composition gives rise to a very rich line spectrum in the ultraviolet.

4. New Mn II energy levels

After adding to the Kurucz line list the missing atomic species relevant for the star and available in the literature, and after comparing the observed spectrum with the synthetic spectrum computed with the abundances given in Table 1, we concentrated on the Mn II lines. Line data for this ion were recently updated and critically evaluated by Kramida & Sansonetti (2013). They list 3969 Mn II lines from 933.0682 Å to 9907.26 Å from which they derived 277 even energy levels and 377 odd energy levels. All the lines are classified as transitions between known energy levels, except for 188 lines observed in the laboratory spectra but not classified. Most of these lines lie in the 2380–2700 Å region. Numerous unidentified lines can be observed in this region in HD 175640. The wavelengths of several of them are the unclassified Mn II lines.

To determine new energy levels we used the same method adopted by Castelli & Kurucz (2010) when they derived new Fe II energy levels from the optical UVES spectrum of HR 6000.

Predicted energy levels and log gf values for Mn II were computed by Kurucz with his version of the Cowan (1981) code (Kurucz 2011). The calculation included 50 even configurations d⁶, d⁵4d–12d, d⁴4s4d–4s9d, d⁵4s–12s, d⁴4s², d⁴4s5s–4s9s, d⁵5g–9g, d⁴4s5g–4s9g, d⁵7i–9i, d⁴4s7i–4s9i, d⁴4s9i, and d⁴4p² with 19686 eigenvalues that were fitted with the least-squares method to 277 known levels. The 41 odd configurations included d⁵4p–10p, d⁴4s4p–4s10p, d³s²4p, d⁵4f–10f, d⁴4s4f–4s10f, d⁵6h–9h, d⁴4s6h–4s9h, d⁵8k–9k, and d⁴4s8k–4s9k with 19820 eigenvalues which were least-squares fitted to 377 known levels. The calculations were done in intermediate coupling with all configuration interactions included, with scaled Hartree-Fock starting guesses, and with Hartree-Fock transition integrals. A total of 5146779 lines were saved from the transition array of which 41882 lines are between known levels and have good wavelengths.

To derive new energy levels for Mn II, we considered predicted wavelengths due to transitions between a measured and a predicted energy level. A predicted line is usually not shifted more than ±10 Å from the corresponding unknown observed line. However, owing to the large number of predicted lines, of unidentified lines, and blends in the ultraviolet spectrum of HD 175640, it is not a straightforward task to find the correct correspondence between a predicted line and an unidentified line observed in the spectrum.

The computed line list was sorted into tables of all the lines with log gf ≥ -3.0 connected to every predicted level. At first, we extracted the lines with log gf ≥ -1.0 so that only the strongest lines were examined. When at least two strong predicted lines originate from the same predicted level, we searched in the spectrum for an unidentified line with wavelength close to the first of the two predicted lines. From the observed wavelength and the known energy level involved in the transition we derived a possible value for the unknown energy level. We used this energy value to derive the corresponding wavelength of the second predicted line. If there is an unidentified line at this position in the observed spectrum we checked the energy value on other predicted lines connected to the examined predicted level. If the test is positive for a sample of lines (usually from 3 to 5), we assign the checked energy to the unknown level. Otherwise, we selected some other observed unidentified line in the spectrum and we repeated the procedure. We proceed in this way until we find that value for the unknown energy that produces lines which are all observed in the spectrum, but not identified.

Whenever one or more new levels was found, the whole semiempirical calculation was repeated to produce improved predicted wavelengths and log gf values.

Using the above procedure we identified 58 new energy levels from the HST-STIS spectrum of HD 175640; five of them have odd parity, the others have even parity. We added 15 more new odd parity energy levels that we obtained from the UVES spectrum of HD 175640. The new energy levels are listed in Table 2 and Table 3.

The search for new energy levels in the optical region was rather unsuccessful. The energies 98423.93 cm⁻¹ and 98424.00 cm⁻¹ of the two levels belonging to the term 3d⁵(⁶S)4f⁷F^o, were obtained from the strong lines of the triplets at 5294 Å and 5295 Å, which are part of the multiplet 3d⁵(⁶S)4d e⁷D^o – 3d⁵(⁶S)4f⁷F^o. All the transitions of this multiplet give rise to five strong Mn II lines observed at 5294, 5295, 5297, 5299, and 5302 Å. Because the lines computed with the energies listed by Kramida & Sansonetti (2013) are more or less displaced from the observed lines, we modified the energy

of the upper level in order to better fit the observed line position. Table 4 compares the energies of the $3d^5(^6S)4f^7F^o$ term adopted in this paper with those listed in the NIST database. The corresponding wavelengths of the multiplet are also listed. Figure 2 shows the five Mn II lines computed with wavelengths from this paper listed in the first column of Table 4 and those from Kramida & Sansonetti (2013) listed in the last column of Table 4.

The energies 113840.6 cm^{-1} , 113840.7 cm^{-1} , and 113840.8 cm^{-1} of the three levels with $J=2, 1$, and 0 respectively, belonging to the odd parity term $3d^5(^6S)6f^7F^o$ were simply extrapolated from the levels with higher J quantum number. The transitions from these levels contribute to two weak and blended lines observed at 7166.4 and 7167.4 Å .

The energy 119197.79 cm^{-1} was assumed for all the levels with J from 0 to 7 of the term $3d^5(^6S)8f^7F^o$. They give rise to two weak lines observed at 5180.271 Å and 5181.649 Å . The first one is blended with Fe II 5180.312 Å .

Finally, the observed lines arising from the three last energy levels of Table 3 are listed in the Table A.1 available online. In addition to the corrections of the odd parity levels shown in Table 4, we modified two other odd parity energy levels from Kramida & Sansonetti (2013) on the basis of all the transitions observed in the spectrum related to them. They are: 81802.746 cm^{-1} instead of 81803.31 cm^{-1} ($3d^5(^2I)4p\ z^1I^o$, $J=6$) and 94230.94 cm^{-1} instead of 94231.20 cm^{-1} ($3d^5(^2S)4p\ z^3P^o$, $J=2$).

5. The new Mn II lines

The new Mn II lines due to transitions from the new Mn II energy levels listed in Table 2 and Table 3 are given in Table A.1, available online. The new Mn II lines are mostly concentrated in the $2380\text{--}2700\text{ Å}$ interval. The upper energy levels (Cols. 1–4) were derived as described in Sect. 4; the lower energy levels (Cols. 5–8) were taken from Kramida & Sansonetti (2013); the wavelength given in Col. 9 is the Ritz wavelength in air for $\lambda \geq 2000\text{ Å}$, in vacuum for $\lambda < 2000\text{ Å}$. The $\log gf$ values (Col. 10) were computed by Kurucz with the semi-empirical method, the wavelengths given in Col. 11 are the observed laboratory wavelengths listed by Kramida & Sansonetti (2013). In addition to the 109 newly classified lines, we give a different assignment for 24 other lines of the Kramida & Sansonetti (2013) tabulation. This implies different energy levels for these 24 lines. The last column of Table A.1 indicates whether the new line is listed in the NIST database without any classification (unassigned), listed with a different classification (DIFFERENT assign.), or not listed at all.

Figure 3 shows the Mn II spectrum in the regions $2383\text{--}2390\text{ Å}$ interval computed before and after the determination of both new energy levels and new Mn II lines. Figure 4 compares the observed spectrum of HD 175640 with the synthetic spectrum computed both with and without the new Mn II lines. The improvement of the comparison between the observed and computed spectra due to the addition of the new Mn II lines is evident.

6. Mn II $\log gf$ values and manganese abundance

For Mn II, Kramida & Sansonetti (2013) tabulate in the NIST database the experimental $\log gf$ values from Den Hartog et al. (2011), Kling & Griesmann (2000), and Kling, Schnabel & Griesmann (2001). For lines with upper levels $3d^5(^6S)4p\ z^5P^o$ and $3d^5(^6S)4p\ z^7P^o$, they adjusted the $\log gf$ values given in the last two papers according to the lifetime value recommended

by Den Hartog et al. (2011) for these levels. Furthermore, they adopted the semi-empirical $\log gf$ values from Kurucz (1988) for several lines.

The comparison of the experimental $\log gf$ values (KGS) with the last version of the semi-empirical oscillator strengths from Kurucz (2014) (K14) is shown in Fig. 5. The average of the difference $\log gf(K14) - \log gf(KGS)$ is $+0.068 \pm 0.397$. The trend is small, but the dispersion around the zero is rather large. However, on a total of 193 lines, 51 of them have a difference in $\log gf$ less than 0.1 dex.

In order to extract a set of lines well suited to deriving the manganese abundance, we presumed that lines with small differences between experimental and semi-empirical $\log gf$ values could be used to this purpose. Table 5 lists 18 Mn II lines with $\log gf$ values from K14 and from either KGS or Den Hartog et al. (2011) differing less than 0.06 dex. The smallest difference (0.004 dex) occurs for the line at 4343.983 Å . This line gives an abundance of -3.80 dex. We checked this abundance on the other lines listed in Table 5. Instead of deriving approximately the same abundance from the lines, we found a variety of values ranging from -3.8 dex to -4.2 dex, as is shown in the last column of Table 5. We investigated whether the abundance differences are related to the uncertainty associated with the experimental $\log gf$ values. According to Kramida & Sansonetti (2013) their accuracy is ≤ 0.009 dex for the score A^+ and decreases up to ≤ 0.24 dex for the score D . If we restrict our analysis to the lines in Table 5 with the scores B , B^+ and A^+ , corresponding to uncertainties not greater than 0.04 dex, we find that the discrepancies in the abundances are not reduced. Figure 6 shows that the problem occurs because some strong Mn II lines have profiles that cannot be reproduced by computations for any adopted abundance. For instance, while the high abundance of -3.8 dex improves the agreement between the observed and computed profiles of the strong line with no wings at 2716.793 Å , it is too high for the broad line at 2949.204 Å , which develops unobserved strong wings for abundances larger than -4.20 dex. We note that the experimental $\log gf$ value of this line is determined with an uncertainty ≤ 0.009 dex, so that it cannot be the cause of the weaker than observed computed core for the -4.20 dex abundance, well suited to fitting the wings. An increase in the microturbulent velocity from 0.0 km sec^{-1} to higher values does not solve the problem because the effect is the same as that due to an increase in the abundance. The difficulty in fitting the profile of several strong lines can be due to missing hyperfine structure (but this is not the case of the line at 2716.793 Å); $\log gf$ uncertainties larger than those estimated; the adopted model, which may have an incorrect stratification in the uppermost layers; and some manganese vertical abundance stratification causing a manganese accumulation in the upper layers.

The analysis of all the Mn II lines in HD 175640 from 1250 Å to 10000 Å , has shown that the most reliable manganese abundance is that derived from the lines of multiplet 1 in the $3438\text{--}3500\text{ Å}$ spectral region. The measured equivalent widths listed in Castelli & Hubrig (2004) together with the $\log gf$ values from Den Hartog et al. (2011) give an average abundance of -4.17 ± 0.03 dex. This abundance reproduces very well both the wings and the core of the lines at 3441.896 , 3460.315 , 3482.904 , and 3488.676 Å . For the other lines of the multiplet the observed core is a little stronger than the computed value.

There are eight points outside the borders of Fig. 5, which were not plotted. They correspond to the lines listed in Table 6 for which the difference between the K14 and KSG $\log gf$ values is larger than ± 1.0 dex. The uncertainties of the experimental




Fig. 2. Improved energy levels of Mn II. Five lines from multiplet $3d^5(^6S)4d e^7D - 3d^5(^6S)4f e^7F^o$ are computed using the wavelengths and the energies from Kramida & Sansonetti (2013) (bottom panel), and the current values (upper panel). The new and old wavelengths and energies are listed in Table 4. The labels are the same as in Fig. 1.

Table 5. Abundances from Mn II lines with experimental and semi-empirical $\log gf$ values that differ less than ± 0.06 dex.

wave Å	$\log gf$ K14	E(low) cm ⁻¹	J(low)	Term	E(up) cm ⁻¹	J(up)	Term	hfs	$\log gf$ KSG	Acc.	$\Delta \log gf$ K14-KSG	Abund
1678.643	-2.681	9472.993	2.0	(⁶ S)4s a ⁵ S	69044.910	2.0	(⁴ P)4p z ³ P	no	-2.638	D ⁺	-0.043	-4.20
2535.659	-0.478	27583.590	4.0	(⁴ G)4s a ⁵ G	67009.217	3.0	(⁴ P)4p z ⁵ D	no	-0.502	B ⁺	+0.024	-3.80
2535.977	-1.024	27588.534	3.0	(⁴ G)4s a ⁵ G	67009.217	3.0	(⁴ P)4p z ⁵ D	no	-1.045	B ⁺	+0.021	-3.80
2543.457	-0.119	27589.360	2.0	(⁴ G)4s a ⁵ G	66894.130	1.0	(⁴ G)4p z ⁵ F	no	-0.138	B	+0.019	-4.20
2557.543	-0.310	27588.534	3.0	(⁴ G)4s a ⁵ G	66676.833	2.0	(⁴ P)4p z ⁵ D	no	-0.346	C ⁺	+0.036	-4.10
2716.793	-0.664	29889.534	3.0	(⁴ P)4s a ⁵ P	66686.739	3.0	(⁴ G)4p z ⁵ F	yes	-0.646	B ⁺	-0.018	-3.80
2762.558	-1.422	32857.270	3.0	(⁴ D)4s b ⁵ D	69044.910	2.0	(⁴ P)4p z ³ P	no	-1.471	D ⁺	+0.049	-4.20
2933.785	-1.421	32818.440	0.0	(⁴ D)4s b ⁵ D	66894.130	1.0	(⁴ G)4p z ⁵ F	yes	-1.458	B	+0.037	-4.20
2935.362	-2.047	32836.740	1.0	(⁴ D)4s b ⁵ D	66894.130	1.0	(⁴ G)4p z ⁵ F	no	-2.002	C ⁺	-0.045	-3.85
2949.204	+0.313	9472.993	2.0	(⁶ S)4s a ⁵ S	43370.537	3.0	(⁶ S)4p z ⁵ P	no	+0.253 ^a	A ⁺	+0.060	-4.20
2955.139	-0.972	32857.270	3.0	(⁴ D)4s b ⁵ D	66686.739	3.0	(⁴ G)4p z ⁵ F	no	-0.987	B ⁺	+0.015	-3.80
2989.730	-1.934	33248.660	4.0	(⁴ G)4s a ³ G	66686.739	3.0	(⁴ G)4p z ⁵ F	no	-1.924	D ⁺	-0.010	-3.90
3050.657	-0.159	36274.620	2.0	(⁴ P)4s b ³ P	69044.910	2.0	(⁴ P)4p z ³ P	no	-0.206	B ⁺	+0.047	-4.15
3146.121	-2.546	34910.770	4.0	d6 b ³ 3G	66686.739	3.0	(⁴ G)4p z ⁵ F	no	-2.523	D	-0.023	-4.25
3204.878	-1.115	37851.490	3.0	d6 a ³ 3D	69044.910	2.0	(⁴ P)4p z ³ P	no	-1.097	B ⁺	-0.018	-4.15
4206.368	-1.584	43528.661	5.0	(⁴ F)4s a ⁵ F	67295.446	4.0	(⁴ P)4p z ⁵ D	yes	-1.553	C ⁺	-0.031	-4.05
4292.233	-1.581	43395.395	3.0	(² D)4s c ³ D	66686.739	3.0	(⁴ G)4p z ⁵ F	yes	-1.544	D ⁺	-0.037	-4.20
4343.983	-1.105	43528.661	5.0	(⁴ F)4s a ⁵ F	66542.539	5.0	(⁴ G)4p z ⁵ F	yes	-1.109	C ⁺	+0.004	-3.80

This value is from Den Hartog et al. (2011)




Fig. 3. Upper panel: Mn II synthetic spectrum for the parameters of HD 175640 ($T_{\text{eff}}=12000$ K, $\log g=3.95$, $v \sin i=2.5$ km $^{-1}$, $[\text{Mn}/\text{H}]=+2.4$) computed with the Mn II line list computed by Kurucz before this work (2009 line list); lower panel: the same, but with the new Mn II lines added in the line list.

Table 6. Mn II lines with experimental and semi-empirical $\log gf$ values that differ more than ± 1.00 dex.

wave Å	$\log gf$ K14	E(low) cm $^{-1}$	J(low)	Term	E(up) cm $^{-1}$	J(up)	Term	hfs	$\log gf$ KSG	Acc.	$\Delta \log gf$ K14-KSG	$\log gf$ for the −4.20 dex abund
2559.416	−0.350	27583.590	4.0	(^4G)4s a ^5G	66643.296	4.0	(^4G)4p z ^5F	no	−1.424	D	+1.074	−0.35
2717.524	−0.895	29889.534	3.0	(^4P)4s a ^5P	66676.833	2.0	(^4P)4p z ^5D	yes	−2.184	E	+1.289	−0.70
2719.736	−0.319	29919.444	2.0	(^4P)4s a ^5P	66676.833	2.0	(^4P)4p z ^5D	yes	−1.500	D	+1.181	−0.319
2845.848	−2.732	31514.710	4.0	d6 a ^3F	66643.296	4.0	(^4G)4p z ^5F	no	−1.274	D $^+$	−1.458	−2.85
2956.005	−1.325	32857.270	3.0	(^4D)4s b ^5D	66676.833	2.0	(^4P)4p z ^5D	no	−3.168	D	+1.843	−0.75
2993.603	−2.421	33147.710	5.0	(^4G)4s a ^3G	66542.539	5.0	(^4G)4p z ^5F	no	−1.061	D $^+$	−1.360	−2.35
4325.047	−2.355	43528.661	5.0	(^4F)4s a ^5F	66643.296	4.0	(^4G)4p z ^5F	yes	−1.115	C $^+$	−1.240	−2.55
4393.385	−2.646	44139.031	1.0	(^2D)4s c ^3D	66894.130	1.0	(^4G)4p z ^5F	yes	−1.458	D $^+$	−1.188	−2.70

$\log gf$ values is high because it is included between ≤ 0.08 dex (score C $^+$) and > 0.24 dex (score E). We note that the source of the $\log gf$ values for the first three lines in Table 6 was erroneously indicated in the NIST database as c88 (Kurucz 1988) instead of T7259 (Kling et al. 2001). The $\log gf$ value needed to reproduce the line profile when the abundance of −4.20 dex is adopted is given in the last column of Table 6. For all the lines, this $\log gf$ value is closer to that from K14 than to that from KGS. This result suggests that, for some lines, the experimental $\log gf$ values may be less reliable than the calculated values.

7. Comparison of the observed and computed spectra

The plot of the comparison of the observed and computed spectra of HD 175640 in the whole 1250–3040 Å region is available at the Castelli website³.

When we compare the whole observed region from 1250 Å to 3046 Å with the synthetic spectrum we are far from satisfied. From 1250 Å to 1692 Å the agreement is poor, except for a few lines. In fact, in spite of the rather high resolving power ($R \sim 40000 - 50000$), the blends are so numerous and formed by so many components that it is not easy to compute them correctly. There are numerous missing lines, lines computed that

³ <http://wwwuser.oats.inaf.it/castelli/hd175640stis/tab1250-3040.html>




Fig. 4. Upper panel: Comparison of the STIS spectrum of HD 175640 (black line) with a synthetic spectrum (red line) computed with a line list including the new Mn II lines; lower panel: the same, but the synthetic spectrum is computed with a line list not including the lines given in Table A.1 available online. Only the Mn II lines are indicated with red labels in the plot. The meaning of the labels is the same as in Fig. 1.

are too strong, lines computed that are too weak, and lines probably affected by small wavelength errors. Furthermore, for a few elements, the abundance is different for the different ionization states. Between 1692 Å and 2332 Å the comparison is even worse because all the above problems are increased by the rather low resolving power ($R \sim 25000\text{--}30000$). Only in the 2332 – 3040 Å region does the comparison show a rather good agreement. This range is less crowded with lines than the previous regions and was observed with a very high resolving power (~ 120000).

7.1. Problematic Mn II lines

If we restrict the discussion only to the Mn II lines, we see that there are several lines with computed profiles without an observed counterpart. For instance, all the lines arising from the even parity level 43131.51 cm^{-1} ($3d^6\text{ a}^1\text{D}, J=2$) are predicted as being strong, but they are either not observed at all or have a very weak observed counterpart. Another remarkable line is that at 2703.213 Å, which is predicted as very strong with both the Kurucz (2014) and Kling et al. (2001) $\log gf$ values, i.e. -0.484 and -0.808 , respectively, but is not observed at all in the spectrum. It would be too long to list here all the Mn II lines computed as too strong compared to the observed ones. In some cases, a

possible explanation for the disagreement is the severe mixing affecting one of the two energy levels involved in the transition and the difficulty in computing it. Furthermore, it is very likely that the computed stellar atmosphere is inadequate to explain all the lines observed in the spectrum, in particular for manganese which is very overabundant. In addition to the shortcoming affecting the core of the strong Mn II lines which, as discussed in Sect. 6, is computed as too weak compared to the observed core, there are several Mn II lines in the optical spectrum that seem to be weakened by some emission. While the computed lines are strong the observed lines are weak or even absent. All these lines are due to transitions between high-energy levels like the blend at 6446.188, 6446.247, 6446.330 Å, the lines at 6462.220, 6462.465, 6462.800, 6463.200, 6463.627 Å, those at 9903.853, 9904.428, 9905.216, 9906.2, and 9907.2 Å, among others.

On the other hand, there are a few strong lines observed in the spectrum that are identified as Mn II in the NIST database, but which are either not predicted by the Kurucz (2014) $\log gf$ values or are predicted as much too weak. Table A.2, available online, lists the most significant lines.




Fig. 5. Comparison of the experimental $\log gf$ values from Kling & Griesmann (2000) and Kling et al. (2001) with the semi-empirical computed values from Kurucz (2014). There are still eight points outside the figure, four with $\log gf > 1.0$ and four with $\log gf < -1.0$.

7.2. Absorption components of resonance lines

A few strong resonance lines are either double with a red component or are very broad with a redshifted line core. In this last case, the profile is the blend of the line predicted at the Ritz wavelength with an unpredicted redshifted line. In Table 7 we list in Col. 2 the Ritz wavelength and in Cols. 3 and 4 the wavelengths λ_{obs} and λ_{comp} of the two components when they can be disentangled. In this case λ_{obs} is the same as the Ritz wavelength, otherwise λ_{obs} is the wavelength of the blend of the two components unless the line is so saturated that the core lies below the zero level of the normalized flux scale and cannot be measured. The velocity shift of the component is given in Col. 5. We estimate an error of about 2.5 km sec^{-1} related to an uncertainty of 0.02 \AA in the position of the component on the wavelength scale. The average velocity shift is $8.29 \pm 0.56 \text{ km s}^{-1}$. We note that the Zn II line at 2062.001 \AA is predicted but not observed, while an unidentified line is present at 2062.060 \AA . We assumed that it is the red component of Zn II 2062.001 \AA . However, it could be the unpredicted line of some other element as well.

Other similar peculiarities observed in HD 175640 are the bump and the broad weak component affecting the red wing of the Ca II profiles at 3933 \AA (K-line) and 3968 \AA (H-line), respectively. Furthermore, strong red components can be observed for the Na I lines at 5890 \AA and 5896 \AA (Castelli & Hubrig, 2004). We measured a velocity shift for the Na I components equal to $+7.4 \pm 0.7 \text{ km sec}^{-1}$ and $+7.7 \pm 0.7 \text{ km sec}^{-1}$, respectively.

We cannot say whether the redshifted components are of interstellar or circumstellar origin.

8. Conclusions

Ultraviolet and optical stellar spectra of the HgMn slowly rotating star HD 175640, observed with both HST-STIS and UVES instruments, were used to extend and discuss the atomic data of Mn II available in the NIST database (Kramida & Sansonetti,

2013). To this purpose, Mn II lines arising both from levels observed in the laboratory and from levels predicted with semi-empirical methods by Kurucz, were adopted.

We assigned wavelengths, energy levels and $\log gf$ values to about 257 Mn II lines. Of them, 109 lines had already been identified as Mn II by Kramida & Sansonetti (2013), but were unclassified. For another 24 lines we assigned different energy levels from those in the NIST database. This implies different energy levels, different $\log gf$ values, and therefore a different intensity for the lines. The new Mn II line data improve the computation of the synthetic spectra of B-type stars, although more work needs to be done on the atomic data, especially in the ultraviolet. For instance, we note that in the NIST database only wavelengths and intensities are listed for the Mn III lines of the $1250\text{--}3040 \text{ \AA}$ region analyzed in the paper. No energy levels and no $\log gf$ values are given for them.

As by-product of this study we have extended to the ultraviolet region the abundance analysis performed by Castelli & Hubrig (2004) in the optical region. Except for cobalt, iridium, and platinum, we confirmed the optical abundances within the error limits of the equivalent widths analysis made by Castelli & Hubrig (2004). Abundances for elements not observed in the visible, such as B, N, Al, Cl, V, Zn, Ge, As, Ag, and Cd were obtained. The abundance pattern is similar to that of other HgMn stars which have underabundances of the light elements and overabundances of some iron group elements and of some heavy elements. In the case of HD 175640, the most overabundant elements are Ti, Cr, Mn, Ga, As, Br, Y, Zr, Rh, Pd, Ag, Xe, Yb, Au, and Hg.

References

- Adelman, S. J. 1994, MNRAS, 266, 97
- Ansbacher, W., Pinnington, E. H., Bahr, J. L., & Kernahan, J. A. 1985, Canadian Journal of Physics, 63, 1330
- Ayres, T. R. 2010, ApJS, 187, 149
- Ayres, T. R. 2014, <http://casa.colorado.edu/~ayres/ASTRAL/>




Fig. 6. The comparison of the observed spectrum (black line) with synthetic spectra computed for a manganese abundance of -4.2 dex (red line) and -3.8 dex (blue line). The plot on the left shows how the computed core of the line at 2716.793 Å increases with increasing abundance, although it does not fit the observed spectrum even with -3.8 dex. A still larger abundance would develop unobserved wings. The line is computed with the hyperfine structure included. The strongest hyperfine components are those at 2716.796 Å and 2716.806 Å. The plot on the right shows how the wings of the strong Mn II line at 2949.204 Å are well fitted by the profile computed with the abundance of -4.2 dex. The higher abundance gives rise to unobserved wings. The meaning of the labels is the same as in Fig. 1.

- Bäckström, E., Nilsson, H., Engström, L., Hartman, H., & Mannervik, S. 2013, *Journal of Physics B Atomic Molecular Physics*, 46, 205001
- Bidelman, W. P. 1962, *AJ*, 67, 111
- Biémont, É., Blagoev, K., Engström, L., et al. 2011, *MNRAS*, 414, 3350
- Biémont, E., Garnir, H. P., Li, Z. S., et al. 2001, *Journal of Physics B Atomic Molecular Physics*, 34, 1869
- Carpenter, K. G., Ayres, T. R., & the ASTRAL Science Teams 2014, *arXiv:1411.1419*
- Castelli, F., & Bonifacio, P. 1990, *A&AS*, 84, 259
- Castelli, F., Cornachin, M., Morossi, C., & Hack, M. 1985, *A&AS*, 59, 1
- Castelli, F., & Hubrig, S. 2004, *A&A*, 425, 263
- Castelli, F., Johansson, S., & Hubrig, S. 2008, *Journal of Physics Conference Series*, 130, 012003
- Castelli, F., & Kurucz, R. L. 2010, *A&A*, 520, AA57
- Castelli, F., Kurucz, R. L., & Hubrig, S. 2009, *A&A*, 508, 401
- Castelli, F., & Parthasarathy, M. 1995, *Astrophysical Applications of Powerful New Databases*, A.S.P. Conf. Ser. 78, 151
- Corliss, C. H., & Bozman, W. R. 1962, *NBS Monograph*, Washington: US Department of Commerce, National Bureau of Standards, —c1962.
- Cowan, R. D. 1981, *The Theory of Atomic Structure and Spectra* (Berkeley: Univ. California Press)
- Cowley, C. R., & Hensberge, H. 1981, *ApJ*, 244, 252
- Den Hartog, E. A., Lawler, J. E., Sobek, J. S., Sneden, C., & Cowan, J. J. 2011, *ApJS*, 194, 35
- Dworetzky, M. M. 1969, *ApJ*, 156, L101
- Holt, R. A., Scholl, T. J., & Rosner, S. D. 1999, *MNRAS*, 306, 107
- Iglesias, L., & Velasco, R. 1964, *The spectrum of the Mn⁺ ion*, by Iglesias, Laura.; Velasco, R. Madrid : Consejo Superior de Investigaciones Cientificas, [1964],
- Ivarsson, S., Wahlgren, G. M., Dai, Z., Lundberg, H., & Leckrone, D. S. 2004, *A&A*, 425, 353
- Kling, R., & Griesmann, U. 2000, *ApJ*, 531, 1173
- Kling, R., Schnabel, R., & Griesmann, U. 2001, *ApJS*, 134, 173
- Kramida, A., & Sansonetti, J. E. 2013, *ApJS*, 205, 14
- Kramida, A., Ralchenko, Y., Reader, J., & NIST ASD Team 2014, *NIST Atomic Spectra Database* (ver. 5.2), available at: <http://physics.nist.gov/asd>
- Kurucz, R. L. 1988, *Trans. IAU, XXB*, edited by McNally (Dordrecht: Kluwer, 1988), pp. 168-172. Data available at <http://kurucz.harvard.edu/linelists/gfall/>
- Kurucz, R. 1993, *SYNTHE Spectrum Synthesis Programs and Line Data*. Kurucz CD-ROM No. 18. Cambridge, Mass.: Smithsonian Astrophysical Observatory, 1993., 18. Data available at <http://kurucz.harvard.edu/linelists/gfall/>
- Kurucz, R. L. 2005, *Memorie della Societa Astronomica Italiana Supplementi*, 8, 14
- Kurucz, R. L. 2011, *Canadian Journal of Physics*, 89, 417. Data available at <http://kurucz.harvard.edu/atoms/>
- Kurucz, R. L. 2014, File *hyper250155.pos* available at <http://kurucz.harvard.edu/atoms/2501/>
- Lundberg, H., Johansson, S. G., Larsson, J., et al. 1996, *ApJ*, 469, 388
- Moore, C. E. 1950, *NBS Circular 488*, Washington: US Government Printing Office (USGPO)

- Nave, G., & Johansson, S. 2013, ApJS, 204, 1
- Nielsen, K. E., Wahlgren, G. M., Proffitt, C. R., Leckrone, D. S., & Adelman, S. J. 2005, AJ, 130, 2312
- Peterson, R. C., & Kurucz, R. L. 2015, ApJS, 216, 1
- Proffitt, C. R., Brage, T., Leckrone, D. S., et al. 1999, ApJ, 512, 942
- Quinet, P. 1996, Phys. Scr., 54, 483
- Quinet, P., Biémont, E., Palmeri, P., et al. 2012, A&A, 537, AA74
- Rosberg, M., & Wyart, J.-F. 1997, Phys. Scr., 55, 690
- Ryabchikova, T. A., Piskunov, N. E., Kupka, F., & Weiss, W. W. 1997, Baltic Astronomy, 6, 244, <http://vald.inasan.ru/~vald3/php/vald.php>
- Ryabtsev, A. N., Wyart, J.-F., Joshi, Y. N., Raassen, A. J. J., & Uylings, P. H. M. 1993, Phys. Scr., 47, 45
- Sugar, J., & Corliss, C. 1985, Washington: American Chemical Society, 1985,
- Uylings, P. H. M., & Raassen, A. J. J. 1997, A&AS, 125, 539
- Warner, B., & Kirkpatrick, R. C. 1969, MNRAS, 142, 265
- Wyart, J.-F., Blaise, J., & Joshi, Y. N. 1995, Phys. Scr., 52, 535
- Wyart, J.-F., Joshi, Y. N., Tchang-Brillet, L., & Raassen, A. J. J. 1996, Phys. Scr., 53, 174

Appendix A: Heavy elements in HD 175640

We based the abundance analysis in the ultraviolet mostly on the lines listed by Castelli et al. (1985) and by Castelli & Bonifacio (1990) in their study of HR 6000 and ι Her, respectively. The multiplet numbers given in the above papers refer to the Ultraviolet Multiplet Table by Moore (1950). In addition, we made a wide use of the NIST database (Kramida et al. 2014) to search for the most intense lines of a given element. In this section we give some more details only about heavy elements from copper to mercury.

Copper (Cu) Z=29: The abundance of -6.50 dex was estimated from the Cu II lines at 1358.773, 1367.951, 1472.395, 2112.100, 2135.981, 2192.268, 2242.618, and 2247.003 Å. All the lines are blended. The log gf values are data computed by Kurucz in 2011 and were taken from his database.

Zinc (Zn) Z=30: The zinc abundance was derived from the line of Zn II at 2064.227 Å. The log gf value $+0.070$ is from the NIST database. The Zn II line at 2062.001 Å originating from a lower level with energy equal to 0.00 cm^{-1} is predicted, but not observed. The log gf value -0.329 is from the Kurucz database. The other line from the zero energy level at 2025.484 Å is computed too weak and is blended with a strong unidentified line that we assumed to be of interstellar or circumstellar origin. No Zn III lines were observed.

Gallium (Ga) Z=31: For Ga I, Ga II, and Ga III, wavelengths, energy levels, and log gf values were taken from the NIST database when available. For Ga I, only the lines at 2874.23, 2943.64, and 2944.17 Å were well observed. For the Ga II lines with log gf values not available in the NIST database, we used log gf values from Nielsen et al. (2005) (lines at 1463.576, 1473.690, 1483.453, and 1483.903 Å). For the remaining lines log gf values were computed from the lifetime measurements of Szbacher et al. (1985) or estimated on the basis of laboratory intensities and excitation energies (Castelli & Parthasarathy 1995). The abundance -5.43 dex ($+3.6$) derived from the optical region adequately reproduces the unblended Ga II lines at 1473.690, 1483.903, 1504.334, 1536.276 Å, as well as the blended lines at 1514.505, 1535.312 Å. The Ga II broad line at 1414.399 Å has too narrow wings, but the classical broadening parameters adopted here are probably not correct. Vice versa, both Ga III profiles at 1495.045 and 1534.462 Å display wings that are too broad. Only these two Ga III lines have log gf values available in the NIST database. For other lines we adopted estimated values. No isotopic and hyperfine structure was considered in the computations.

Germanium (Ge) Z=32: Several Ge II lines were observed.

From the line at 1649.19 Å we derived an underabundance of 1.7 dex relative to the solar value. We adopted $\log gf = -0.28$ from the NIST database

Arsenic (As) Z=33: The lines of As II at 1263.77, 1266.34, 1280.987, and 1287.54 Å were observed in the spectrum. We adopted log gf values from Warner & Kirkpatrick (1969) for all them. The observed lines are adequately predicted for an overabundance of -7.50 dex.

Yttrium (Y) Z=39: For the Y II lines, we used the Kurucz database, which includes hyperfine components for several lines. All the lines are blended, except for the strong line at 2422.18 Å. The abundance of -6.66 dex ($+3.2$) derived from the optical spectrum is well suited to reproducing the observed profile. We added to the line list the Y III lines from Biémont et al. (2011). We modified the two wavelengths 2817.037 Å and 2946.01 Å in 2817.027 Å and 2945.995 Å, in order to match the observed spectrum. The Y III lines at 2367.228, 2414.643, 2817.027, and 2945.995 Å were observed. They are strong lines, either unblended or marginally blended. The abundance from Y III is larger than the abundance from Y II by about 1.0 dex.

Zirconium (Zr) Z=40: Several weak Zr III lines were observed in the spectrum, but no Zr II line. A few Zr III lines are unblended, like that at 2102.283 Å. The Zr III line data from the NIST database were adopted. The Zr III abundance agrees, within the error limits, with the value we obtained from the Zr II optical lines.

Rhodium (Rh) Z=45: Numerous weak Rh II lines were observed in the spectrum. The line at 1604.45 Å is unblended. Oscillator strengths from Bäckström et al. (2013), Quinet et al. (2012), and Corliss & Bozman (1962) were used. The wavelengths from the NIST database agree more closely with the stellar wavelengths than the wavelengths adopted by Bäckström et al. (2013). The difference is on the order of 0.02 Å. The abundance of -8.50 dex, which was estimated from the UVES spectra on the basis of estimated log gf values, reproduces the ultraviolet lines in a satisfactory way. None of the Rh III lines listed in the NIST database was identified.

Palladium (Pd) Z=46: Numerous Pd II lines were identified. All the lines from the $4d^8 5s-4d^8 5p$ transitions and the strongest lines from the $4d^8 5p-4d^8 6s$ transitions, together with their oscillator strengths, from Quinet (1996) were added in the line lists. Additional lines were taken from Lundberg et al. (1996) and from the NIST database. Unblended lines are those at 2351.347, 2367.966, 2388.310, 2414.7303, 2426.867, 2433.102, 2446.713, 2457.257, 2472.502, 2486.256, 2488.914, 2505.729, 2565.505, 2569.544, 2635.93, and 2658.72 Å. None of the Pd I lines listed in the NIST database was identified. For Pd III, only the line at 1782.55 Å, which according to the NIST data is the one with the largest intensity in the 1250-3040 Å range, perhaps contributes to a strong blend observed at 1782.6 Å.

Silver (Ag) Z=47: The Ag II lines at 2246.412, 2248.749, 2357.917, and 2411.345 Å were observed. The log gf values from the NIST database were used when available, otherwise we used the Corliss & Bozman (1962) data, as we did for the line at 2438.325 Å, which is computed as too strong.

Cadmium (Cd) Z=48: The Cd II lines at 2144.393 Å and 2265.019 Å were observed and predicted by adopting a 0.8 overabundance over the solar value. The atomic data are from the Kurucz database. The log gf values are the same as in the NIST database.

Indium (In) Z=49: The In II line at 1586.331 Å, which is predicted as rather strong for solar abundance and log gf value from the NIST database, is the main component of a complex

blend formed by several other lines. We were unable to determine the indium abundance from the blend.

Barium (Ba) Z=56: The two strongest Ba II lines at 2304.249 Å and 2335.267 Å are heavily blended, and so they cannot be used to confirm the abundance derived from the optical region. No other Ba II lines were observed.

Ytterbium (Yb) Z=70: We adopted the abundance of -8.10 dex ($[+3.0]$) derived from the UVES spectra. We investigated only the Yb II lines with $\log gf$ available in the NIST database. The lines at 2185.716, 2653.745, and 2750.478 can be observed as single weak lines. Some other lines are weak components of blends. We added to the line list all the Yb III lines from Biémont et al. (2001) with an upper energy level lower than 100000 cm^{-1} . There are numerous Yb III lines in the spectrum. Their profile is very sharp and would require an abundance about 1 dex larger than that derived from the Yb II lines in order to agree with the computed profiles. The same behavior for Yb II and Yb III abundances was observed in the visible (Castelli & Hubrig 2004).

Osmium (Os) Z=76: No Os II lines were observed

Iridium (Ir) Z=77: Except for the weak line observed at 2245.750 Å, no other Ir II lines can be conclusively identified in the spectrum. From this line we derived an abundance of -11.15 dex, i.e. an underabundance of $[-0.5]$. The $\log gf$ values for Ir II were taken from the Kurucz database. None of the Ir II lines listed in Ivarsson et al. (2004) was observed in the spectrum, even assuming a solar abundance. The Ir II line at 3042.553 Å is a minor contributor of a blend with Ti II. The blending was not considered when we analyzed the UVES spectrum, so that we derived a solar iridium abundance (Castelli & Hubrig 2004).

Platinum (Pt) Z=78: For Pt II and Pt III we adopted the lines and the transition probabilities listed by Wyart et al. (1995) and by Ryabtsev et al. (1993), respectively. We did not find any clear evidence for a Pt overabundance. Only the Pt II line at 1777.086 Å is predicted with solar abundance and observed. It is a blend with Fe II 1777.058 Å, which is computed as weaker than observed, but an overabundance of platinum gives rise to computed lines not observed in the spectrum. We believe that the line at 4514.124 Å, identified as Pt II in the UVES spectrum (Castelli & Hubrig 2004), is actually due to some other element.

Gold (Au) Z=79: Gold is overabundant by 3.6 dex. The abundance from the optical region adequately reproduces the numerous Au II lines observed in the ultraviolet. A few of them are unblended, good examples being the Au II lines at 1469.142, 1673.587, 1740.475, 1793.297 (blend), and 1800.579 Å. There are several predicted Au III lines, but they are all blended, except for those at 1365.382 and 1385.768 Å. We adopted the Au II and Au III line lists from Rosberg & Wyart (1997) and Wyart et al. (1996), respectively.

Mercury (Hg) Z=80: The $\log gf$ values for Hg I and Hg II were taken from the NIST database. For Hg III we considered only the three lines at 1360.509, 1647.482, and 1738.540 Å listed by Profitt et al. (1999). The abundance of -6.60 dex is well suited to reproducing the lines in the ultraviolet, while -6.30 dex derived from the Hg II line at 3984 Å (Castelli & Hubrig 2004) is too large. We did not consider any hyperfine structure in the computation of the ultraviolet lines. For Hg I, the line at 1849.499 Å is blended, while the line at 2536.521 Å is very weak. No other lines with available $\log gf$ values were observed. Numerous lines of Hg II were observed and computed. For instance, the line at 1942.273 Å is strong, but it is blended with Mn II at 1942.344 Å. Two lines at 1321.712 Å and

1331.738 Å are single and well reproduced, those at 1354.289, 1539.142, and 1869.226 Å are strong, but blended. For Hg III, the line at 1360.509 Å is computed as too strong, that at 1647.471 Å would be adequately reproduced if the Mn III line at 1647.497 Å were omitted; finally, the line at 1738.540 Å is blended with numerous other components.

Table 1. Abundances $\log(N_{elem}/N_{tot})$ for HD 175640 [12000K,3.95] adopted to compute the ultraviolet synthetic spectrum. Solar abundances are from Asplund et al. (2009).

Elem.	HD175640		Sun
	UV	Visible	
He I		-1.73	-1.11
Be II		-10.64	-10.99
B II	-8.75		-9.34
C I	< -4.00	-4.11±0.23	-3.61
C II	-4.00	-4.05±0.16	-3.61
N I	-5.95	≤ -5.78	-4.21
O I	-3.28	-3.18±0.11	-3.35
Ne I		-4.35	-4.11
Na I		-5.47	-5.80
Mg I	-4.69	-4.64 ± 0.06	-4.44
Mg II	-4.69	-4.71 ± 0.07	-4.44
Al I		< -7.50	-5.59
Al II	-7.00		-5.59
Al III	-7.00		-5.59
Si I	-4.80		-4.53
Si II	-4.80	-4.72 ± 0.08	-4.53
Si III	-4.80	-4.58 ± 0.04	-4.53
Si IV	-4.80		-4.53
P I	< -6.28		-6.63
P II	-6.28	-6.28±0.08	-6.63
P III	-6.28		-6.63
S I	-5.30		-4.92
S II	-5.30	-5.12±0.03	-4.92
Cl I	-7.50		-6.54
Ca I		-5.26	-5.70
Ca II	-5.54	-5.67 ± 0.25	-5.70
Sc II	-9.08	-9.08 ± 0.15	-8.89
Sc III	-9.08		-8.89
Ti II	-5.67	-5.67±0.11	-7.09
Ti III	-5.67		-7.09
V II	-9.94	≤ -9.04	-8.11
Cr I		-5.22 ± 0.09	-6.40
Cr II	-5.36	-5.41 ± 0.07	-6.40
Cr III	-5.36		-6.40
Mn I	-4.20	-4.20±0.08	-6.61
Mn II	-4.20	-4.25 ± 0.04	-6.61
Mn III	-4.20:		-6.61
Fe I	-4.83	-4.78±0.08	-4.54
Fe II	-4.83	-4.84±0.13	-4.54
Fe III	-4.83		-4.54
Co II	-9.00	-8.08:	-7.05
Ni II	-6.09	-6.09±0.16	-5.82
Ni III	-6.09		-5.82
Cu I		-6.52	-7.85
Cu II	-6.50		-7.85
Zn II	-8.70		-7.48
Ga I	-5.43		-9.00
Ga II	-5.43	-5.43±0.04	-9.00
Ga III	< -5.43 ?		-9.00
Ge II	-10.1		-8.39
As II	-7.50		-9.74
Br II		-7.12±0.04	-9.50
Sr II		-8.41	-9.17
Y II	-6.66	-6.66±0.20	-9.83
Y III	> -6.66		-9.83

Table 1. cont.

elem	HD175640		Sun
	UV	Visible	
Zr II		-8.67±0.17	-9.46
Zr III	-8.50		-9.46
Rh II	-8.50	-8.50	-11.13
Pd I		-6.41 ± 0.30	-10.47
Pd II	-6.30		-10.47
Pd III	-6.30		-10.47
Ag II	-8.50		-11.10
Cd II	-9.53		-10.33
Xe II		-5.96 ± 0.20	-9.80
Ba II		-9.27	-9.86
Pr III		-9.62	-11.32
Nd III		-9.57 ± 0.08	-10.62
Yb II	-8.10	-8.10 ± 0.19	-11.20
Yb III	> -8.10	-7.31±0.01	-11.20
Os II		-10.55	-10.64
Ir II	-11.15	-10.66:	-10.66
Pt	-10.42	-7.63	-10.42
Au II	-7.51	-7.51±0.06	-11.12
Au III	-7.51		-11.12
Hg I	-6.60	-6.19 ± 0.18	-10.87
Hg II	-6.60	-6.53 ± 0.23	-10.87
Hg III	-6.60		-10.87

Table 2. New energy levels of Mn II from the STIS-HST spectrum.

Designation	J	Energy cm ⁻¹
3d ⁴ (³ D)4s4p(³ P)	⁷ F ^o 3.0	82303.415
3d ⁴ (³ D)4s4p(³ P)	⁷ F ^o 4.0	82564.493
3d ⁵ (b ² D)4p	³ F ^o 2.0	100320.586
	4.0	100725.176
3d ⁵ (b ² D)4p	³ P ^o 1.0	101807.469
3d ⁶ (⁴ G)4d	⁵ F 1.0	106351.582
	2.0	106348.454
	3.0	106340.485
	4.0	106324.758
	5.0	106298.960
3d ⁵ (⁴ G)4d	⁵ G 2.0	106408.011
	3.0	106408.348
	4.0	106406.642
	5.0	106400.766
	6.0	106387.310
3d ⁵ (⁴ G)4d	³ D 3.0	107192.929
3d ⁵ (⁴ G)4d	³ I 5.0	107155.877
	6.0	107145.038
	7.0	107114.165
3d ⁵ (⁴ G)4d	³ G 3.0	107999.208
	4.0	108007.750
	5.0	108006.293
3d ⁵ (⁴ G)4d	³ F 2.0	108420.629
	3.0	108425.105
	4.0	108420.479
3d ⁶ (⁴ P)4d	⁵ P 1.0	108617.023
	3.0	108399.148
3d ⁶ (⁴ P)4d	⁵ F 1.0	108511.206
	2.0	108521.377
	3.0	108561.169
	4.0	108604.283
	5.0	108661.385
3d ⁵ (⁴ G)4d	³ H 4.0	108907.450
	5.0	108906.437
	6.0	108896.070
3d ⁶ (⁴ P)4d	³ D 1.0	109709.489
	2.0	109572.155
	3.0	109370.762
3d ⁵ (⁴ P)4d	³ F 2.0	110664.620
	3.0	110611.014
3d ⁵ (⁴ P)4d	³ P 2.0	110998.198
3d ⁵ (⁴ D)4d	⁵ G 2.0	111744.523
	3.0	111753.977
	4.0	111761.546
	5.0	111761.008
	6.0	111741.583

Table 2. Cont.

Designation	J	Energy cm ⁻¹
3d ⁵ (⁴ D)4d	⁵ P 3.0	111831.748
3d ⁵ (⁴ D)4d	⁵ S 2.0	111991.133
3d ⁵ (⁴ D)4d	⁵ F 2.0	112141.748
	3.0	112106.537
	4.0	112046.403
	5.0	111943.809
3d ⁵ (⁴ D)4d	⁵ D 4.0	113199.572
3d ⁵ (² I)4d	³ I 5.0	118544.936
	6.0	118574.338
	7.0	118585.339
3d ⁵ (² I)4d	¹ K 7.0	119152.797
3d ⁵ (² I)4d	¹ G 4.0	119937.245

Table 3. New energy levels of Mn II from the optical region (UVES spectrum).

Designation	J	Energy cm ⁻¹
3d ⁵ (⁶ S)4f	⁷ F ^o 1.0	98423.93
	0.0	98424.00
3d ⁵ (⁶ S)6f	⁷ F ^o 2.0	113840.6
	1.0	113840.7
	0.0	113840.8
3d ⁵ (⁶ S)8f	⁷ F ^o 6.0	119197.79
	5.0	119197.79
	4.0	119197.79
	3.0	119197.79
	2.0	119197.79
	1.0	119197.79
3d ⁵ (⁴ G)4f	⁵ H ^o 7.0	125218.640
	9.0	125243.338
3d ⁵ (⁴ G)4f	⁵ I ^o 6.0	125257.344

Table 4. The multiplet $3d^5(^6S)4d\,e^7D - 3d^5(^6S)4f\,e^7F^o$. The energies and wavelengths from this paper are compared with those from Kramida & Sansonetti (2013). We note that with the new levels, all 15 of the possible lines of the $^7D-^7F$ multiplet are known.

$\lambda(\text{CKC})^a$ Å	El(KS) ^b cm ⁻¹	Eu(CKC) ^a cm ⁻¹	Jl,Ju	Eu(KS) ^b cm ⁻¹	$\lambda(\text{KS})^b$ Å
5294.276	79540.93	98424.00	1,0	---	---
5294.296	79540.93	98423.93	1,1	---	---
5294.318	79540.93	98423.858	1,2	98423.63	5294.379
5295.356	79544.71	98423.93	2,1		
5295.376	79544.71	98423.858	2,2	98423.63	5295.439
5295.400	79544.71	98423.773	2,3	98423.63	5295.439
5297.003	79550.50	98423.858	3,2	98423.63	5297.063
5297.024	79550.50	98423.773	3,3	98423.63	5297.063
5297.068	79550.50	98423.624	3,4	98423.60	5297.072
5299.288	79558.56	98423.773	4,3	98423.63	5299.327
5299.329	79558.56	98423.624	4,4	98423.60	5299.336
5299.387	79558.56	98423.424	4,5	98423.59	5299.338
5302.327	79569.22	98423.624	5,4	98423.60	5302.331
5302.382	79569.22	98423.424	5,5	98423.59	5302.333
5302.421	79569.22	98423.281	5,6	98423.56	5302.342

^a CKC: this paper;

^b KS: Kramida & Sansonetti (2013)

Table 7. Lines affected by a redshifted component.

Elem	$\lambda(\text{Ritz})$ (Å)	$\lambda(\text{obs})$ (Å)	λ_{comp} (Å)	V_{shift} km s ⁻¹	Notes
Mg I	2852.126	2852.126	2852.20	7.78	double
Mg II	2795.528				saturated
	2802.705				saturated
Si II	1808.013	1808.03			
S II	1250.584	1250.60			
Mn II	2576.104	2576.104	2576.18	8.84	double
	2593.721	2593.721	2593.80	9.13	double
	2605.680	2605.68	2605.75	8.05	double
Fe II	2343.495	2343.53			
	2373.735	2373.735	2373.80	8.21	double
	2382.037				saturated
	2585.876	2585.876	2585.94	7.42	double
	2599.395				saturated
Zn II	2025.484	2025.51			
	2062.001	--	2062.06	8.57	

Table A.1. New energy levels of Mn II and the predicted lines observed in the spectrum.

Upper level				Lower level				$\lambda(\text{Ritz})$	$\log gf$	$\lambda(\text{NIST})$	Notes
Eup(cm ⁻¹)	Conf.	Term	J	Elow(cm ⁻¹)	Conf.	Term	J	Å	K14	Å	
82303.415	3d ⁴ (⁵ D)4s4p(³ P)	⁷ F ^o	3.0	14325.866	3d ⁶	a ⁵ D	4.0	1471.074	-3.902		
				14593.835	3d ⁶	a ⁵ D	3.0	1476.896	-3.407		
				14781.205	3d ⁶	a ⁵ D	2.0	1480.994	-1.901		
				29869.420	3d ⁶	a ³ P	2.0	1907.160	-3.174		
				43395.395	3d ⁵ (² D)4s	c ³ D	3.0	2569.394	-0.806		
				43696.217	3d ⁵ (⁴ F)4s	a ⁵ F	3.0	2589.416	-2.294	2589.42	unassigned
				43850.341	3d ⁵ (⁴ F)4s	a ⁵ F	2.0	2599.795	-1.673	2599.794	unassigned
				44521.540	3d ⁵ (² F)4s	b ³ F	4.0	2645.984	-1.855	2645.99	unassigned
				46903.320	3d ⁵ (² D)4s	b ¹ D	2.0	2824.020	-0.594		
82564.493	3d ⁴ (⁵ D)4s4p(³ P)	⁷ F ^o	4.0	32857.270	3d ⁵ (⁴ D)4s	b ⁵ D	3.0	2011.131	-2.962		
				43395.395	3d ⁵ (² D)4s	c ³ D	3.0	2552.269	-2.421	2552.272	unassigned
				43528.661	3d ⁵ (⁴ F)4s	a ⁵ F	5.0	2560.983	-1.875	2560.981	unassigned
				43537.186	3d ⁵ (⁴ F)4s	a ⁵ F	4.0	2561.542	-1.716	2561.542	unassigned
				43696.217	3d ⁵ (⁴ F)4s	a ⁵ F	3.0	2572.024	-2.444	2572.023	unassigned
100320.586	3d ⁵ (b ² D)4p	³ F ^o	2.0	31761.210	3d ⁶	a ³ F	2.0	1458.590	-1.756		
				33278.730	3d ⁵ (⁴ G)4s	a ³ G	3.0	1491.605	-1.533		
				35004.790	3d ⁶	b ³ G	3.0	1531.023	-1.231		
				37811.880	3d ⁶	a ³ D	1.0	1599.777	-1.533		
				62564.771	3d ⁵ (² D)4s	d ³ D	1.0	2647.810	-0.092	2647.811	DIFFERENT assign.
				62572.199	3d ⁵ (² D)4s	d ³ D	2.0	2648.331	-0.277	2648.339	unassigned
				62587.547	3d ⁵ (² D)4s	d ³ D	3.0	2649.408	-1.196		
100725.176	3d ⁵ (b ² D)4p	³ F ^o	4.0	31514.710	3d ⁶	a ³ F	4.0	1444.868	-1.594		
				33147.710	3d ⁵ (⁴ G)4s	a ³ G	5.0	1479.783	-1.133		
				34762.098	3d ⁶	b ³ G	5.0	1516.000	-1.053		
				37851.490	3d ⁶	a ³ D	3.0	1590.490	-1.376		
				49517.690	3d ⁵ (² G)4s	c ³ G	5.0	1952.839	-1.587		
				52383.780	3d ⁵ (² F)4s	d ³ F	4.0	2067.960	-1.386		
				62587.547	3d ⁵ (² D)4s	d ³ D	3.0	2621.300	+0.408	2621.300	unassigned
101807.469	3d ⁵ (b ² D)4p	³ P ^o	1.0	62564.771	3d ⁵ (² D)4s	d ³ D	1.0	2547.480	-0.526	2547.48	unassigned
				62572.199	3d ⁵ (² D)4s	d ³ D	2.0	2547.962	-0.338	2547.966	unassigned
106351.582	3d ⁶ (⁴ G)4d	⁵ F	1.0	64456.720	3d ⁵ (⁴ G)4p	z ⁵ G ^o	2.0	2386.200	-0.175	2386.206	unassigned
				66676.833	3d ⁵ (⁴ P)4p	z ⁵ D ^o	2.0	2519.737	-0.870		
				66894.130	3d ⁵ (⁴ G)4p	z ⁵ F ^o	1.0	2533.614	-0.211	2533.610	unassigned
				66901.494	3d ⁵ (⁴ G)4p	z ⁵ F ^o	2.0	2534.087	-0.725		
				70150.760	3d ⁵ (⁴ P)4p	y ⁵ F ^o	1.0	2761.552	-1.333		
106348.454	3d ⁶ (⁴ G)4d	⁵ F	2.0	64456.720	3d ⁵ (⁴ G)4p	z ⁵ G ^o	2.0	2386.378	-0.595		
				64473.421	3d ⁵ (⁴ G)4p	z ⁵ G ^o	3.0	2387.330	-0.094		
				66645.116	3d ⁵ (⁴ P)4p	z ⁵ D ^o	1.0	2517.922	-1.189		
				66676.833	3d ⁵ (⁴ P)4p	z ⁵ D ^o	2.0	2519.935	-0.488		
				66686.739	3d ⁵ (⁴ G)4p	z ⁵ F ^o	3.0	2520.565	-0.443		
				66894.130	3d ⁵ (⁴ G)4p	z ⁵ F ^o	1.0	2533.815	-0.440	2533.817	unassigned.
				66901.494	3d ⁵ (⁴ G)4p	z ⁵ F ^o	2.0	2534.288	-0.312		
				67009.217	3d ⁵ (⁴ P)4p	z ⁵ D ^o	3.0	2541.228	-0.989		
106340.485	3d ⁶ (⁴ G)4d	⁵ F	3.0	64473.421	3d ⁵ (⁴ G)4p	z ⁵ G ^o	3.0	2387.784	-0.494		
				64494.140	3d ⁵ (⁴ G)4p	z ⁵ G ^o	4.0	2388.967	+0.032		
				66643.296	3d ⁵ (⁴ G)4p	z ⁵ F ^o	4.0	2518.312	-0.342	2518.311	DIFFERENT assign.
				66676.833	3d ⁵ (⁴ P)4p	z ⁵ D ^o	2.0	2520.442	-0.519		
				66686.739	3d ⁵ (⁴ G)4p	z ⁵ F ^o	3.0	2521.071	-0.003	2521.081	unassigned
				66901.494	3d ⁵ (⁴ G)4p	z ⁵ F ^o	2.0	2534.800	-0.465	2534.804	unassigned
				67009.217	3d ⁵ (⁴ P)4p	z ⁵ D ^o	3.0	2541.743	-0.511	2541.740	DIFFERENT assign.

Table A.1. Cont.

Upper level				Lower level				$\lambda(\text{Ritz})$	$\log gf$	$\lambda(\text{NIST})$	Notes
Eup(cm ⁻¹)	Conf.	Term	J	Elow(cm ⁻¹)	Conf.	Term	J	Å	K14	Å	
106324.758	3d ⁶ (⁴ G)4d	⁵ F	4.0	64494.140	3d ⁵ (⁴ G)4p	z ⁵ G ^o	4.0	2389.865	−0.611		
				64518.890	3d ⁵ (⁴ G)4p	z ⁵ G ^o	5.0	2391.280	+0.178	2391.281	unassigned
				65658.650	3d ⁵ (⁴ G)4p	z ⁵ H ^o	5.0	2458.306	−0.937		
				66542.539	3d ⁵ (⁴ G)4p	z ⁵ F ^o	5.0	2512.929	−0.462	2512.932	unassigned
				66643.296	3d ⁵ (⁴ G)4p	z ⁵ F ^o	4.0	2519.310	+0.320	2519.312	DIFFERENT assign.
				66686.739	3d ⁵ (⁴ G)4p	z ⁵ F ^o	3.0	2522.072	−0.278		
				67009.217	3d ⁵ (⁴ P)4p	z ⁵ D ^o	3.0	2542.760	−0.951		
				67295.446	3d ⁵ (⁴ P)4p	z ⁵ D ^o	4.0	2561.409	−0.930	2561.412	unassigned
				70497.790	3d ⁵ (⁴ D)4p	y ⁵ F ^o	4.0	2790.371	−0.953		
106298.960	3d ⁵ (⁴ G)4d	⁵ F	5.0	64518.890	3d ⁵ (⁴ G)4p	z ⁵ G ^o	5.0	2392.757	−0.993		
				64550.040	3d ⁵ (⁴ G)4p	z ⁵ G ^o	6.0	2394.542	+0.303	2394.54	unassigned
				65754.820	3d ⁵ (⁴ G)4p	z ⁵ H ^o	6.0	2465.702	−1.037		
				66542.539	3d ⁵ (⁴ G)4p	z ⁵ F ^o	5.0	2514.560	+0.558	2514.567	unassigned
				66643.296	3d ⁵ (⁴ G)4p	z ⁵ F ^o	4.0	2520.949	−0.398	2520.958	unassigned.
				70657.595	3d ⁵ (⁴ D)4p	y ⁵ F ^o	5.0	2804.902	−0.782		
106408.011	3d ⁵ (⁴ G)4d	⁵ G	2.0	64456.720	3d ⁵ (⁴ G)4p	z ⁵ G ^o	2.0	2382.990	+0.182	2382.991	DIFFERENT assign
				64473.421	3d ⁵ (⁴ G)4p	z ⁵ G ^o	3.0	2383.939	−0.267		
				65483.120	3d ⁵ (⁴ G)4p	z ⁵ H ^o	3.0	2442.760	−0.781		
				66894.130	3d ⁵ (⁴ G)4p	z ⁵ F ^o	1.0	2529.996	−0.415	2529.999	unassigned
				66901.494	3d ⁵ (⁴ G)4p	z ⁵ F ^o	2.0	2530.467	−0.999		
106408.348	3d ⁵ (⁴ G)4d	⁵ G	3.0	64456.720	3d ⁵ (⁴ G)4p	z ⁵ G ^o	2.0	2382.971	−0.565		
				64473.421	3d ⁵ (⁴ G)4p	z ⁵ G ^o	3.0	2383.920	+0.229	2383.921	unassigned
				64494.140	3d ⁵ (⁴ G)4p	z ⁵ G ^o	4.0	2385.099	−0.087	2385.096	unassigned
				65483.120	3d ⁵ (⁴ G)4p	z ⁵ H ^o	3.0	2442.740	−0.775		
				65566.030	3d ⁵ (⁴ G)4p	z ⁵ H ^o	4.0	2447.699	−0.799	2447.697	unassigned
				66676.833	3d ⁵ (⁴ P)4p	z ⁵ D ^o	2.0	2516.136	−0.552		
				66686.739	3d ⁵ (⁴ G)4p	z ⁵ F ^o	3.0	2516.764	−0.705		
106406.642	3d ⁵ (⁴ G)4d	⁵ G	4.0	66901.494	3d ⁵ (⁴ G)4p	z ⁵ F ^o	2.0	2530.446	−0.446	2530.447	unassigned
				64473.421	3d ⁵ (⁴ G)4p	z ⁵ G ^o	3.0	2384.017	−0.404		
				64494.140	3d ⁵ (⁴ G)4p	z ⁵ G ^o	4.0	2385.196	+0.344		
				64518.890	3d ⁵ (⁴ G)4p	z ⁵ G ^o	5.0	2386.605	−0.093	2386.605	DIFFERENT assign.
				65566.030	3d ⁵ (⁴ G)4p	z ⁵ H ^o	4.0	2447.802	−0.474	2447.803	unassigned
				65658.650	3d ⁵ (⁴ G)4p	z ⁵ H ^o	5.0	2453.366	−0.682	2453.371	unassigned
				66643.296	3d ⁵ (⁴ G)4p	z ⁵ F ^o	4.0	2514.122	−0.713		
				66686.739	3d ⁵ (⁴ G)4p	z ⁵ F ^o	3.0	2516.872	−0.103	2516.875	unassigned
106400.766	3d ⁵ (⁴ G)4d	⁵ G	5.0	67009.217	3d ⁵ (⁴ P)4p	z ⁵ D ^o	3.0	2537.475	−0.703	2537.480	unassigned
				64494.140	3d ⁵ (⁴ G)4p	z ⁵ G ^o	4.0	2385.530	−0.387	2385.532	unassigned
				64518.890	3d ⁵ (⁴ G)4p	z ⁵ G ^o	5.0	2386.940	+0.479	2386.940	unassigned
				64550.040	3d ⁵ (⁴ G)4p	z ⁵ G ^o	6.0	2388.717	−0.289	2388.72	DIFFERENT assign.
				65658.650	3d ⁵ (⁴ G)4p	z ⁵ H ^o	5.0	2453.720	−0.319	2453.721	unassigned
				65754.820	3d ⁵ (⁴ G)4p	z ⁵ H ^o	6.0	2459.526	−0.466	2459.53	DIFFERENT assign
				66542.539	3d ⁵ (⁴ G)4p	z ⁵ F ^o	5.0	2508.137	−1.088		
				66643.296	3d ⁵ (⁴ G)4p	z ⁵ F ^o	4.0	2514.494	+0.145	2514.503	unassigned
106387.310	3d ⁵ (⁴ G)4d	⁵ G	6.0	64518.890	3d ⁵ (⁴ G)4p	z ⁵ G ^o	5.0	2387.707	−0.513		
				64550.040	3d ⁵ (⁴ G)4p	z ⁵ G ^o	6.0	2389.485	+0.616	2389.49	unassigned
				65754.820	3d ⁵ (⁴ G)4p	z ⁵ H ^o	6.0	2460.340	−0.319		
				65847.030	3d ⁵ (⁴ G)4p	z ⁵ H ^o	7.0	2465.937	−0.199	2465.944	unassigned.
				66542.539	3d ⁵ (⁴ G)4p	z ⁵ F ^o	5.0	2508.984	+0.281	2508.984	unassigned
107192.929	3d ⁵ (⁴ G)4d	³ D	3.0	66643.296	3d ⁵ (⁴ G)4p	z ⁵ F ^o	4.0	2465.368	−0.954		
				67812.087	3d ⁵ (⁴ G)4p	z ³ F ^o	3.0	2538.543	−0.477	2538.543	unassigned
				67865.945	3d ⁵ (⁴ G)4p	z ³ F ^o	4.0	2542.020	+0.499	2542.024	unassigned
				73395.444	3d ⁵ (⁴ D)4p	y ³ D ^o	3.0	2957.936	−0.752	2957.920	DIFFERENT assign.

Table A.1. Cont.

Upper level				Lower level				$\lambda(\text{Ritz})$	$\log gf$	$\lambda(\text{NIST})$	Notes
Eup(cm ⁻¹)	Conf.	Term	J	Elow(cm ⁻¹)	Conf.	Term	J	Å	K14	Å	
107155.877	3d ⁵ (⁴ G)4d	³ I	5.0	67846.240	3d ⁵ (⁴ G)4p	z ³ H ^o	5.0	2543.142	−0.656		
				67910.560	3d ⁵ (⁴ G)4p	z ³ H ^o	4.0	2547.310	+0.810	2547.31	unassigned
107145.038	3d ⁵ (⁴ G)4d	³ I	6.0	67744.440	3d ⁵ (⁴ G)4p	z ³ H ^o	6.0	2537.27	−0.644	2537.265	unassigned
				67846.240	3d ⁵ (⁴ G)4p	z ³ H ^o	5.0	2543.843	+0.885	2543.851	unassigned
107114.165	3d ⁵ (⁴ G)4d	³ I	7.0	64550.040	3d ⁵ (⁴ G)4p	z ³ G ^o	6.0	2348.677	−1.321		
				67744.440	3d ⁵ (⁴ G)4p	z ³ H ^o	6.0	2539.260	+0.960	2539.272	unassigned
107999.208	3d ⁵ (⁴ G)4d	³ G	3.0	67766.817	3d ⁵ (⁴ G)4p	z ³ F ^o	2.0	2484.809	+0.024	2484.808	unassigned
				67910.560	3d ⁵ (⁴ G)4p	z ³ H ^o	4.0	2493.720	−0.379	2493.716	unassigned
				70518.06	3d ⁵ (⁴ G)4p	z ³ G ^o	3.0	2667.215	+0.351	2667.215	unassigned
108007.750	3d ⁵ (⁴ G)4d	³ G	4.0	67812.087	3d ⁵ (⁴ G)4p	z ³ F ^o	3.0	2487.080	+0.149	2487.089	unassigned
				67846.240	3d ⁵ (⁴ G)4p	z ³ H ^o	5.0	2489.195	−0.283	2489.206	unassigned
				70518.06	3d ⁵ (⁴ G)4p	z ³ G ^o	3.0	2666.607	−0.996		
				70546.340	3d ⁵ (⁴ G)4p	z ³ G ^o	4.0	2668.620	+0.440	2668.620	unassigned.
108006.293	3d ⁵ (⁴ G)4d	³ G	5.0	67744.440	3d ⁵ (⁴ G)4p	z ³ H ^o	6.0	2482.991	−0.181		
				67865.945	3d ⁵ (⁴ G)4p	z ³ F ^o	4.0	2490.508	+0.234	2490.508	unassigned
				70527.61	3d ⁵ (⁴ G)4p	z ³ G ^o	5.0	2667.390	+0.553	2667.393	unassigned
				73683.440	3d ⁵ (⁴ D)4p	y ³ F ^o	4.0	2912.658	−0.989		
108420.629	3d ⁵ (⁴ G)4d	³ F	2.0	67766.817	3d ⁵ (⁴ G)4p	z ³ F ^o	2.0	2459.050	+0.081	2459.042	unassigned
				67812.087	3d ⁵ (⁴ G)4p	z ³ F ^o	3.0	2461.791	−0.783		
				70518.06	3d ⁵ (⁴ G)4p	z ³ G ^o	3.0	2637.558	−0.066		
				71078.501	3d ⁵ (⁴ P)4p	z ³ D ^o	1.0	2677.145	−0.832		
108425.105	3d ⁵ (⁴ G)4d	³ F	3.0	67812.087	3d ⁵ (⁴ G)4p	z ³ F ^o	3.0	2461.520	+0.210	2461.525	DIFFERENT assign.
				67865.945	3d ⁵ (⁴ G)4p	z ³ F ^o	4.0	2464.789	−0.771		
				70497.790	3d ⁵ (⁴ D)4p	y ³ F ^o	4.0	2645.837	−1.420	2635.84	unassigned
				70546.340	3d ⁵ (⁴ G)4p	z ³ G ^o	4.0	2639.215	+0.082	2639.220	unassigned.
				70940.568	3d ⁵ (⁴ P)4p	z ³ D ^o	2.0	2666.974	−0.824		
				73396.334	3d ⁵ (⁴ D)4p	y ³ D ^o	2.0	2853.958	−0.911		
108420.479	3d ⁵ (⁴ G)4d	³ F	4.0	67865.945	3d ⁵ (⁴ G)4p	z ³ F ^o	4.0	2465.070	+0.389	2465.075	unassigned
				70527.61	3d ⁵ (⁴ G)4p	z ³ G ^o	5.0	2638.233	+0.200		
				70745.387	3d ⁵ (⁴ P)4p	z ³ D ^o	3.0	2653.484	−0.753		
				73395.444	3d ⁵ (⁴ D)4p	y ³ D ^o	3.0	2854.262	−0.745		
				73683.440	3d ⁵ (⁴ D)4p	y ³ F ^o	4.0	2877.927	−0.892		
108617.023	3d ⁶ (⁴ P)4d	⁵ P	1.0	66901.494	3d ⁵ (⁴ G)4p	z ⁵ F ^o	2.0	2396.459	−1.205	2396.467	unassigned
				66929.517	3d ⁵ (⁴ P)4p	z ⁵ S ^o	2.0	2398.070	+0.018	2398.082	unassigned
				68417.697	3d ⁵ (⁴ P)4p	y ⁵ P ^o	2.0	2486.853	−0.456		
				68496.694	3d ⁵ (⁴ P)4p	y ⁵ P ^o	1.0	2491.750	−1.034		
				69044.910	3d ⁵ (⁴ P)4p	z ³ P ^o	2.0	2526.272	−1.004	2526.287	unassigned
108399.148	3d ⁶ (⁴ P)4d	⁵ P	3.0	66929.517	3d ⁵ (⁴ P)4p	z ⁵ S ^o	2.0	2410.670	+0.286		
				68284.664	3d ⁵ (⁴ P)4p	y ⁵ P ^o	3.0	2492.113	+0.107	2492.129	DIFFERENT assign.
				68417.697	3d ⁵ (⁴ P)4p	y ⁵ P ^o	2.0	2500.406	−0.255	2500.418	unassigned
				72011.050	3d ⁵ (⁴ D)4p	y ⁵ D ^o	4.0	2747.339	−0.857		
108511.206	3d ⁶ (⁴ P)4d	⁵ F	1.0	66625.296	3d ⁵ (⁴ P)4p	z ⁵ D ^o	0.0	2386.710	−0.141	2386.715	DIFFERENT assign.
				66645.116	3d ⁵ (⁴ P)4p	z ⁵ D ^o	1.0	2387.840	−0.132		
				66676.833	3d ⁵ (⁴ P)4p	z ⁵ D ^o	2.0	2389.651	−0.974		

Table A.1. Cont.

Upper level				Lower level				$\lambda(\text{Ritz})$	$\log gf$	$\lambda(\text{NIST})$	Notes
Eup(cm ⁻¹)	Conf.	Term	J	Elow(cm ⁻¹)	Conf.	Term	J	Å	K14	Å	
108521.377	3d ⁶ (⁴ P)4d	⁵ F	2.0	66645.116	3d ⁵ (⁴ P)4p	z ³ D ^o	1.0	2387.260	+0.036	2387.268	unassigned
				66676.833	3d ⁵ (⁴ P)4p	z ⁵ D ^o	2.0	2389.070	-0.152	2389.075	unassigned
				66894.130	3d ⁵ (⁴ G)4p	z ⁵ F ^o	1.0	2401.542	-0.773		
				66901.494	3d ⁵ (⁴ G)4p	z ⁵ F ^o	2.0	2401.967	-0.278		
				67009.217	3d ⁵ (⁴ P)4p	z ⁵ D ^o	3.0	2408.200	-0.818		
108561.169	3d ⁵ (⁴ P)4d	⁵ F	3.0	72247.730	3d ⁵ (⁴ D)4p	y ⁵ D ^o	3.0	2756.008	-1.771	2756.01	unassigned
				66676.833	3d ⁵ (⁴ P)4p	z ⁵ D ^o	2.0	2386.800	+0.095		
				66686.739	3d ⁵ (⁴ P)4p	z ⁵ F ^o	3.0	2387.364	-0.542		
				66901.494	3d ⁵ (⁴ G)4p	z ⁵ F ^o	2.0	2399.672	-0.012	2399.685	unassigned
				66929.517	3d ⁵ (⁴ P)4p	z ⁵ S ^o	2.0	2401.288	-1.439	2401.298	unassigned
108604.283	3d ⁵ (⁴ P)4d	⁵ F	4.0	67009.217	3d ⁵ (⁴ P)4p	z ⁵ D ^o	3.0	2405.894	+0.032	2405.90	unassigned
				72307.230	3d ⁵ (⁴ D)4p	y ⁵ D ^o	2.0	2757.506	-1.033	2757.51	unassigned
				66686.739	3d ⁵ (⁴ P)4p	z ⁵ F ^o	3.0	2384.909	-0.151	2384.912	unassigned
				67009.217	3d ⁵ (⁴ P)4p	z ⁵ D ^o	3.0	2403.400	+0.445	2403.409	unassigned
				67295.446	3d ⁵ (⁴ P)4p	z ⁵ D ^o	4.0	2420.054	+0.025	2420.06	unassigned
108661.385	3d ⁵ (⁴ P)4d	⁵ F	5.0	72011.050	3d ⁵ (⁴ D)4p	y ⁵ D ^o	4.0	2731.937	-0.934		
				72247.730	3d ⁵ (⁴ D)4p	y ⁵ D ^o	3.0	2749.723	-0.814		
				66643.296	3d ⁵ (⁴ G)4p	z ⁵ F ^o	4.0	2379.201	-0.631		
				67295.446	3d ⁵ (⁴ P)4p	z ⁵ D ^o	4.0	2416.713	+0.750	2416.712	unassigned
				72011.050	3d ⁵ (⁴ D)4p	y ⁵ D ^o	4.0	2727.680	-0.286		
108907.450	3d ⁵ (⁴ G)4d	³ H	4.0	67910.560	3d ⁵ (⁴ G)4p	z ³ H ^o	4.0	2438.470	+0.281	2438.478	unassigned
				70518.06	3d ⁵ (⁴ G)4p	z ³ G ^o	3.0	2604.108	+0.433	2604.124	unassigned
				70546.340	3d ⁵ (⁴ G)4p	z ³ G ^o	4.0	2606.028	-0.870		
108906.437	3d ⁵ (⁴ G)4d	³ H	5.0	67846.240	3d ⁵ (⁴ G)4p	z ³ H ^o	5.0	2434.710	+0.358	2434.676	DIFFERENT assign.
				70497.790	3d ⁵ (⁴ D)4p	y ⁵ F ^o	4.0	2602.803	-0.961		
				70527.610	3d ⁵ (⁴ G)4p	z ³ G ^o	5.0	2604.825	-0.913		
				70546.340	3d ⁵ (⁴ G)4p	z ³ G ^o	4.0	2606.097	+0.525		
108896.070	3d ⁵ (⁴ G)4d	³ H	6.0	67744.440	3d ⁵ (⁴ G)4p	z ³ H ^o	6.0	2429.300	+0.446	2429.297	DIFFERENT assign.
				70527.610	3d ⁵ (⁴ G)4p	z ³ G ^o	5.0	2605.529	+0.621		
109709.489	3d ⁶ (⁴ P)4d	³ D	1.0	69216.059	3d ⁵ (⁴ P)4p	z ³ P ^o	1.0	2468.790	-0.388	2468.786	unassigned.
				69319.330	3d ⁵ (⁴ P)4p	z ³ P ^o	0.0	2475.103	-0.172	2475.095	unassigned
				70940.568	3d ⁵ (⁴ P)4p	z ³ D ^o	2.0	2578.614	-0.996		
				71078.501	3d ⁵ (⁴ P)4p	z ³ D ^o	1.0	2587.821	-0.480		
109572.155	3d ⁶ (⁴ P)4d	³ D	2.0	69044.910	3d ⁵ (⁴ P)4p	z ³ P ^o	2.0	2466.730	-0.480	2466.723	unassigned
				69216.059	3d ⁵ (⁴ P)4p	z ³ P ^o	1.0	2477.192	+0.178	2477.185	unassigned
				70940.568	3d ⁵ (⁴ P)4p	z ³ D ^o	2.0	2587.781	-0.273	2587.76	unassigned
109370.762	3d ⁶ (⁴ P)4d	³ D	3.0	68417.697	3d ⁵ (⁴ P)4p	y ³ P ^o	2.0	2441.080	-0.964		
				69044.910	3d ⁵ (⁴ P)4p	z ³ P ^o	2.0	2479.050	+0.413	2479.05	unassigned
				70342.936	3d ⁵ (⁴ D)4p	y ⁵ F ^o	3.0	2561.506	-0.847		
				70745.387	3d ⁵ (⁴ P)4p	z ³ D ^o	3.0	2588.197	-0.067	2588.217	DIFFERENT assign.
110664.620	3d ⁵ (⁴ P)4d	³ F	2.0	67766.817	3d ⁵ (⁴ G)4p	z ³ F ^o	2.0	2330.407	-0.950		
				70150.760	3d ⁵ (⁴ D)4p	y ⁵ F ^o	1.0	2467.545	-0.400	2467.546	unassigned
				70342.936	3d ⁵ (⁴ D)4p	y ⁵ F ^o	3.0	2482.607	-1.111	2482.604	unassigned
				70940.568	3d ⁵ (⁴ P)4p	z ³ D ^o	2.0	2516.609	-0.584		
				71078.501	3d ⁵ (⁴ P)4p	z ³ D ^o	1.0	2525.378	+0.238		

Table A.1. Cont.

Upper level				Lower level				$\lambda(\text{Ritz})$	$\log gf$	$\lambda(\text{NIST})$	Notes
Eup(cm ⁻¹)	Conf.	Term	J	Elow(cm ⁻¹)	Conf.	Term	J	Å	K14	Å	
110611.014	3d ⁵ (⁴ P)4d	³ F	3.0	67812.087	3d ⁵ (⁴ G)4p	z ³ F ^o	3.0	2335.791	−0.884		
				70231.468	3d ⁵ (⁴ D)4p	y ⁵ F ^o	2.0	2475.753	−0.248	2475.751	unassigned
				70745.387	3d ⁵ (⁴ P)4p	z ³ D ^o	3.0	2507.671	−0.620		
				70940.568	3d ⁵ (⁴ P)4p	z ³ D ^o	2.0	2520.010	+0.412	2520.011	unassigned
110998.198	3d ⁵ (⁴ P)4d	³ P	2.0	69044.910	3d ⁵ (⁴ P)4p	z ³ P ^o	2.0	2382.877	−0.094	2382.88	DIFFERENT assign.
				69216.059	3d ⁵ (⁴ P)4p	z ³ P ^o	1.0	2392.638	−0.666		
				70745.387	3d ⁵ (⁴ P)4p	z ³ D ^o	3.0	2483.549	−1.102	2483.541	unassigned
				71264.390	3d ⁵ (⁴ D)4p	x ⁵ P ^o	1.0	2515.991	−1.079	2515.994	unassigned
				73911.570	3d ⁵ (⁴ P)4p	z ³ S ^o	1.0	2695.590	+0.092	2695.59	DIFFERENT assign.
111744.523	3d ⁵ (⁴ D)4d	⁵ G	2.0	70150.760	3d ⁵ (⁴ D)4p	y ⁵ F ^o	1.0	2403.475	+0.275	2403.477	unassigned
				70231.468	3d ⁵ (⁴ D)4p	y ⁵ F ^o	2.0	2408.148	−0.357	2408.157	unassigned
				71078.501	3d ⁵ (⁴ P)4p	z ³ D ^o	1.0	2458.311	−0.400	2458.32	DIFFERENT assign.
111753.977	3d ⁵ (⁴ D)4d	⁵ G	3.0	70231.468	3d ⁵ (⁴ D)4p	y ⁵ F ^o	2.0	2407.600	+0.430	2407.608	unassigned
				70342.936	3d ⁵ (⁴ D)4p	y ⁵ F ^o	3.0	2414.081	−0.257	2414.08	unassigned
				70745.387	3d ⁵ (⁴ P)4p	z ³ D ^o	3.0	2437.774	−0.872		
				70940.568	3d ⁵ (⁴ P)4p	z ³ D ^o	2.0	2449.433	−0.272	2449.438	unassigned
111761.546	3d ⁵ (⁴ D)4d	⁵ G	4.0	66686.739	3d ⁵ (⁴ P)4p	z ⁵ F ^o	3.0	2217.844	−0.914		
				70342.936	3d ⁵ (⁴ D)4p	y ⁵ F ^o	3.0	2413.640	+0.540	2413.64	unassigned
				70497.790	3d ⁵ (⁴ D)4p	y ⁵ F ^o	4.0	2422.698	−0.191	2422.71	unassigned
				70518.06	3d ⁵ (⁴ G)4p	z ³ G ^o	3.0	2423.889	−0.879		
				70745.387	3d ⁵ (⁴ P)4p	z ³ D ^o	3.0	2437.324	−0.078		
111761.008	3d ⁵ (⁴ D)4d	⁵ G	5.0	66643.296	3d ⁵ (⁴ P)4p	z ⁵ F ^o	4.0	2215.734	−0.735		
				70497.790	3d ⁵ (⁴ D)4p	y ⁵ F ^o	4.0	2422.730	+0.758	2422.749	unassigned
				70546.340	3d ⁵ (⁴ G)4p	z ³ G ^o	4.0	2425.584	−0.764		
				70657.595	3d ⁵ (⁴ D)4p	y ⁵ F ^o	5.0	2432.150	−0.441		
111741.583	3d ⁵ (⁴ D)4d	⁵ G	6.0	66542.539	3d ⁵ (⁴ G)4p	z ⁵ F ^o	5.0	2211.747	−0.649		
				70527.610	3d ⁵ (⁴ G)4p	z ³ G ^o	5.0	2425.625	−0.996		
				70657.595	3d ⁵ (⁴ D)4p	y ⁵ F ^o	5.0	2433.300	+0.868	2433.31	unassigned
111831.748	3d ⁵ (⁴ D)4d	⁵ P	3.0	66929.517	3d ⁵ (⁴ P)4p	z ⁵ S ^o	2.0	2226.368	−0.976		
				67295.446	3d ⁵ (⁴ P)4p	z ⁵ D ^o	4.0	2244.663	−0.689		
				71323.600	3d ⁵ (⁴ D)4p	x ⁵ P ^o	2.0	2467.893	−0.182		
				71390.488	3d ⁵ (⁴ D)4p	x ⁵ P ^o	3.0	2471.975	+0.099	2471.979	unassigned
				72011.050	3d ⁵ (⁴ D)4p	y ⁵ D ^o	4.0	2510.501	−0.210		
				72247.730	3d ⁵ (⁴ D)4p	y ⁵ P ^o	3.0	2525.531	−0.099		
				72307.230	3d ⁵ (⁴ D)4p	y ⁵ D ^o	2.0	2529.315	−0.594	2529.317	unassigned
111991.133	3d ⁵ (⁴ D)4d	⁵ S	2.0	68284.664	3d ⁵ (⁴ P)4p	y ⁵ P ^o	3.0	2287.285	−0.759		
				71264.390	3d ⁵ (⁴ D)4p	x ⁵ P ^o	1.0	2454.646	−0.550	2454.646	unassigned
				71323.600	3d ⁵ (⁴ D)4p	x ⁵ P ^o	2.0	2458.220	−0.212	2458.229	unassigned
				71390.488	3d ⁵ (⁴ D)4p	x ⁵ P ^o	3.0	2462.270	+0.107	2462.276	unassigned
				72247.730	3d ⁵ (⁴ D)4p	y ⁵ P ^o	3.0	2515.386	−0.662		
112141.748	3d ⁵ (⁴ D)4d	⁵ F	2.0	70231.468	3d ⁵ (⁴ D)4p	y ⁵ F ^o	2.0	2385.320	−0.382		
				70342.936	3d ⁵ (⁴ D)4p	y ⁵ F ^o	3.0	2391.682	−0.712		
				70940.568	3d ⁵ (⁴ P)4p	z ³ D ^o	2.0	2426.376	−0.925		
				72307.230	3d ⁵ (⁴ D)4p	y ⁵ D ^o	2.0	2509.627	−0.166		
				72321.020	3d ⁵ (⁴ D)4p	y ⁵ D ^o	1.0	2510.497	+0.051	2510.494	unassigned

Table A.1. Cont.

Upper level				Lower level				$\lambda(\text{Ritz})$	$\log gf$	$\lambda(\text{NIST})$	Notes
Eup(cm ⁻¹)	Conf.	Term	J	Elow(cm ⁻¹)	Conf.	Term	J	Å	K14	Å	
112106.537	3d ⁵ (⁴ D)4d	⁵ F	3.0	70231.468	3d ⁵ (⁴ D)4p	y ⁵ F ^o	2.0	2387.328	-0.921		
				70342.936	3d ⁵ (⁴ D)4p	y ⁵ F ^o	3.0	2393.700	-0.202	2393.70	unassigned.
				70497.790	3d ⁵ (⁴ D)4p	y ⁵ F ^o	4.0	2402.610	-0.596		
				70745.387	3d ⁵ (⁴ P)4p	z ³ D ^o	3.0	2416.993	-0.679		
				71323.600	3d ⁵ (⁴ D)4p	x ⁵ P ^o	2.0	2451.264	-0.746		
				72247.730	3d ⁵ (⁴ D)4p	y ⁵ P ^o	3.0	2508.100	-0.171	2508.102	unassigned
				72307.230	3d ⁵ (⁴ D)4p	y ⁵ D ^o	2.0	2511.850	+0.244	2511.855	unassigned.
112046.403	3d ⁵ (⁴ D)4d	⁵ F	4.0	70342.936	3d ⁵ (⁴ D)4p	y ⁵ F ^o	3.0	2397.152	-0.995		
				70497.790	3d ⁵ (⁴ D)4p	y ⁵ F ^o	4.0	2406.087	+0.102	2406.101	DIFFERENT assign.
				70657.595	3d ⁵ (⁴ D)4p	y ⁵ F ^o	5.0	2415.378	-0.752		
				71390.488	3d ⁵ (⁴ D)4p	x ⁵ P ^o	3.0	2458.923	-0.388	2458.935	unassigned
				72011.050	3d ⁵ (⁴ D)4p	y ⁵ D ^o	4.0	2497.039	-0.336	2497.08	DIFFERENT assign.
				72247.730	3d ⁵ (⁴ D)4p	y ⁵ P ^o	3.0	2511.890	+0.405	2511.905	unassigned
111943.809	3d ⁵ (⁴ D)4d	⁵ F	5.0	67295.446	3d ⁵ (⁴ P)4p	z ⁵ D ^o	4.0	2239.029	-0.788		
				70657.595	3d ⁵ (⁴ D)4p	y ⁵ F ^o	5.0	2421.381	+0.286	2421.38	unassigned
				72011.050	3d ⁵ (⁴ D)4p	y ⁵ D ^o	4.0	2503.455	+0.615	2503.459	unassigned
113199.572	3d ⁵ (⁴ D)4d	⁵ D	4.0	70657.595	3d ⁵ (⁴ D)4p	y ⁵ F ^o	5.0	2349.900	-0.739		
				71390.488	3d ⁵ (⁴ D)4p	x ⁵ P ^o	3.0	2391.096	-0.502		
				72011.050	3d ⁵ (⁴ D)4p	y ⁵ D ^o	4.0	2427.124	+0.005		
				72247.730	3d ⁵ (⁴ D)4p	y ⁵ P ^o	3.0	2441.153	-0.207	2441.147	DIFFERENT assign.
118544.936	3d ⁵ (² I)4d	³ I	5.0	77842.130	3d ⁵ (² I)4p	z ³ K ^o	6.0	2456.090	-0.055		
				78085.260	3d ⁵ (² I)4p	z ³ I ^o	5.0	2470.850	+0.621	2470.849	unassigned
				79800.610	3d ⁵ (² I)4p	y ³ H ^o	4.0	2580.251	+0.111		
				81802.746	3d ⁵ (² I)4p	z ¹ I ^o	6.0	2720.861	-0.935		
				85447.760	3d ⁵ (² H)4p	y ³ I ^o	5.0	3020.527	-0.969		
118574.338	3d ⁵ (² I)4d	³ I	6.0	77842.130	3d ⁵ (² I)4p	z ³ K ^o	6.0	2454.317	-0.040		
				77945.750	3d ⁵ (² I)4p	z ³ K ^o	7.0	2460.577	-0.005	2460.59	DIFFERENT assign.
				78340.740	3d ⁵ (² I)4p	z ³ I ^o	6.0	2484.735	+0.603	2484.735	unassigned
				79112.620	3d ⁵ (² I)4p	z ¹ H ^o	5.0	2533.340	-0.994		
				79739.950	3d ⁵ (² I)4p	y ³ H ^o	5.0	2574.266	+0.208		
118585.339	3d ⁵ (² I)4d	³ I	7.0	85636.170	3d ⁵ (² H)4p	y ³ I ^o	6.0	3035.108	-0.901		
				77820.280	3d ⁵ (² I)4p	z ³ K ^o	8.0	2452.339	+0.012		
				77945.750	3d ⁵ (² I)4p	z ³ K ^o	7.0	2459.910	+0.181		
				78475.470	3d ⁵ (² I)4p	z ³ I ^o	7.0	2492.400	+0.629	2492.404	DIFFERENT assign.
				79592.420	3d ⁵ (² I)4p	y ³ H ^o	6.0	2563.800	+0.316		
119152.797	3d ⁵ (² I)4d	¹ K	7.0	77945.750	3d ⁵ (² I)4p	z ³ K ^o	7.0	2426.033	-0.824		
				79147.220	3d ⁵ (² I)4p	z ¹ K ^o	7.0	2498.898	+0.581		
				81802.746	3d ⁵ (² I)4p	z ¹ I ^o	6.0	2676.577	+0.700		
119937.245	3d ⁵ (² I)4d	¹ G	4.0	79112.620	3d ⁵ (² I)4p	z ¹ H ^o	5.0	2448.76	+0.491	2448.766	DIFFERENT assign.
				79739.950	3d ⁵ (² I)4p	y ³ H ^o	5.0	2486.979	-0.437		
				79800.610	3d ⁵ (² I)4p	y ³ H ^o	4.0	2490.738	-0.327		
				81220.63	3d ⁵ (a ² D)4p	z ¹ F ^o	3.0	2582.098	-0.861		

Table A.2. Lines from the NIST database in the 2400–3040 Å interval which are observed in the spectrum, but are either not predicted or predicted too weak.

$\lambda(\text{NIST})$	$\log gf$	Elow	J	Config.	Term	Eup	J	Config.	Term	Notes
Å		cm ⁻¹				cm ⁻¹				
2409.045	-3.275	55116.333	2	3d ⁴ 4s ²	c ⁵ D	96613.930	2	3d ⁴ (⁵ D)4s4p(³ P ^o)	s ³ F ^o	obs
2421.089	-3.595	49291.360	3	3d ⁶	b ¹ F	90582.550	2	3d ⁴ (³ D)4s4p(³ P ^o)	w ³ F ^o	obs at 2421.10
2431.323	-6.436	49465.150	3	3d ⁵ (² G)4s	c ³ G	90582.550	2	d ⁴ (⁵ D)4s4p(³ P ^o)	w ⁵ F ^o	obs at 2431.33
2441.226	-5.624	73395.444	3	3d ⁵ (⁴ D)4p	y ³ D ^o	114346.060	3	3d ⁵ (⁶ S)7d	⁷ D	obs
2443.392	-5.162	35004.790	3	3d ⁶	b ³ G	75919.107	2	3d ⁶ (⁴ D)4p	y ³ P ^o	obs at 2443.41
2443.809	-4.828	27589.360	2	3d ⁵ (⁴ G)4s	a ⁵ G	68496.694	1	3d ⁵ (⁴ P)4p	y ⁵ P ^o	obs blend
2448.538	-4.389	27589.360	2	3d ⁵ (⁴ G)4s	a ⁵ G	68417.697	2	3d ⁵ (⁴ P)4p	y ⁵ P ^o	obs weak
2458.869	-4.327	29889.534	3	3d ⁵ (⁴ P)4s	a ⁵ P	70546.340	4	3d ⁵ (⁴ G)4p	z ³ G ^o	obs weak
2464.565	-4.819	40759.620	0	3d ⁶	a ¹ S	81322.460	1	3d ⁵ (a ² D)4p	x ³ P ^o	obs blend
2468.395	-4.488	73395.444	3	3d ⁵ (⁴ D)4p	y ³ D ^o	113895.350	2	3d ⁵ (⁶ S)8s	⁵ S	obs at 2468.41
2477.954	-2.323	66542.539	5	3d ⁵ (⁴ G)4p	z ⁵ F ^o	106886.230	4	3d ⁵ (⁴ D)5s	g ⁵ D	obs at 2477.96
2481.228	-2.843	67812.087	3	3d ⁵ (⁴ G)4p	z ³ F ^o	108102.540	3	3d ⁵ (⁴ D)5s	e ³ D	obs blend
2496.885	-4.119	66929.517	2	3d ⁵ (⁴ P)4p	z ⁵ S ^o	106967.350	2	3d ⁵ (⁴ D)5s	e ³ D	obs blend
2500.480	-2.155	66542.539	5	3d ⁵ (⁴ G)4p	z ⁵ F ^o	106522.810	6	3d ⁵ (⁴ G)4d	e ⁵ I	obs
2502.709	-4.230	43311.324	1	3d ⁵ (⁴ F)4s	a ⁵ F	83255.990	2	3d ⁴ (⁵ D)4s4p(³ P ^o)	y ⁷ P ^o	obs blend
2505.093	-3.921	44521.540	4	3d ⁵ (² F)4s	b ³ F	84428.190	5	3d ⁵ (a ² G)4p	x ³ H ^o	obs blend
2518.582	-7.851	79558.560	4	3d ⁵ (⁶ S)4d	e ⁷ D	119251.500	3	3d ⁵ (⁶ S)8f	⁵ F ^o	obs at 2518.56
2518.988	-7.659	79569.220	5	3d ⁵ (⁶ S)4d	e ⁷ D	119255.760	4	3d ⁵ (⁶ S)8f	⁵ F ^o	obs weak
2519.176	-6.675	79569.220	5	3d ⁵ (⁶ S)4d	e ⁷ D	119252.800	5	3d ⁵ (⁶ S)8f	⁵ F ^o	obs at 2519.16
2523.188	-3.091	38720.260	6	3d ⁶	a ¹ I	78340.740	6	3d ⁵ (² I)4p	z ³ I ^o	STRONG
2532.316	-2.138	66686.739	3	3d ⁵ (⁴ G)4p	z ⁵ F ^o	106164.410	4	3d ⁵ (⁴ G)4d	e ⁵ H	obs at 2532.341
2532.320	-2.627	79800.610	4	3d ⁵ (² I)4p	y ³ H ^o	119278.220	5	3d ⁵ (⁶ S)8g	⁵ G	obs at 2532.341
2533.461	-3.086	32787.900	4	3d ⁵ (⁴ D)4s	b ⁵ D	72247.730	3	3d ⁵ (⁴ D)4p	y ⁵ D ^o	STRONG, pred weak
2555.114	-4.599	29919.444	2	3d ⁵ (⁴ P)4s	a ⁵ P	69044.910	2	3d ⁵ (⁴ P)4p	z ³ P ^o	obs blend
2560.172	-4.096	43339.452	2	3d ⁵ (² D)4s	c ³ D	82387.620	3	3d ⁵ (² F)4p	y ³ G ^o	pred weak
2564.447	-3.612	31514.710	4	3d ⁶	a ³ F	70497.790	4	3d ⁵ (⁴ D)4p	y ⁵ F ^o	pred weak
2592.397	-4.652	43537.186	4	3d ⁵ (⁴ F)4s	a ⁵ F	82099.990	4	3d ⁵ (² F)4p	y ⁵ G ^o	STRONG, pred weak
2596.836	-2.244	43696.217	3	3d ⁵ (⁴ F)4s	a ⁵ F	82193.110	2	3d ⁵ (⁴ F)4p	y ⁵ G ^o	STRONG, pred weak
2602.356	-3.726	29869.420	2	3d ⁶	a ³ P	68284.664	3	3d ⁵ (⁴ P)4p	y ⁵ P ^o	pred weak
2639.047	-3.491	43339.452	2	3d ⁵ (² D)4s	c ³ D	81220.630	3	3d ⁵ (a ² D)4p	z ¹ F ^o	obs
2639.706	-3.992	37848.225	2	3d ⁶	a ³ D	75719.955	1	3d ⁵ (⁴ D)4p	y ³ P ^o	STRONG
2666.393	-2.659	72648.860	4	3d ⁵ (² G)4s	d ¹ G	110141.560	4	3d ⁴ (a ³ P)4s4p(³ P ^o)	r ³ F ^o	STRONG
2667.844	-3.398	44521.540	4	3d ⁵ (² F)4s	b ³ F	81993.850	4	3d ⁵ (⁴ F)4p	x ⁵ F ^o	STRONG, blend
2669.604	-2.382	44745.501	2	3d ⁵ (² F)4s	b ³ F	82193.110	2	3d ⁵ (⁴ F)4p	y ⁵ G ^o	STRONG, pred weak
2688.221	-0.785	52383.780	4	3d ⁵ (² F)4s	d ³ F	89572.060	3	3d ⁵ (b ² F)4p	t ³ F ^o	STRONG, pred weak
2716.039	-3.205	29869.420	2	3d ⁶	a ³ P	66676.833	2	3d ⁵ (⁴ P)4p	z ⁵ D ^o	pred weak
2718.382	-3.816	29869.420	2	3d ⁶	a ³ P	66645.116	1	3d ⁵ (⁴ P)4p	z ⁵ D ^o	pred weak
2725.555	-1.652	52383.780	4	3d ⁵ (² F)4s	d ³ F	89062.690	5	3d ⁵ (a ² G)4p	¹ H ^o	pred weak
2736.423	-4.455	49427.371	4	3d ⁵ (² G)4s	c ³ G	85960.622	3	3d ⁵ (⁶ S)5p	x ⁷ P ^o	obs
2739.851	-3.969	49465.150	3	3d ⁵ (² G)4s	c ³ G	85952.690	3.	3d ⁵ (⁴ F)4p	³ F ^o	obs
2760.916	-5.129	30684.960	1	3d ⁶	a ³ P	66894.130	1	3d ⁵ (⁴ G)4p	z ⁵ F ^o	obs weak
2764.089	-2.518	46903.320	2	3d ⁵ (² D)4s	b ¹ D	83070.920	2	3d ⁵ (a ² F)4p	w ³ D ^o	pred weak
2777.168	-4.723	47073.640	3	3d ⁵ (² F)4s	a ¹ F	83070.920	2	3d ⁵ (a ² F)4p	w ³ D ^o	STRONG
2834.640	-6.234	31662.040	3	3d ⁶	a ³ F	66929.517	2	3d ⁵ (⁴ P)4p	z ⁵ S ^o	obs weak
2842.634	-4.902	31761.210	2	3d ⁶	a ³ F	66929.517	2	3d ⁵ (⁴ P)4p	z ⁵ S ^o	obs weak
2844.249	-3.820	46903.320	2	3d ⁵ (² D)4s	b ¹ D	82051.650	3	3d ⁵ (⁴ F)4p	x ⁵ F ^o	pred weak
2844.554	-7.311	74560.181	3	3d ⁵ (⁶ S)5s	e ⁷ 7S	109704.740	2	3d ⁴ (a ³ P)4s4p(³ P ^o)	y ⁵ S ^o	obs at 2844.54
2870.804	-2.714	55759.322	3	3d ⁵ (² F)4s	c ¹ F	90582.550	2	3d ⁴ (³ D)4s4p(³ P ^o)	w ³ F ^o	STRONG
2879.924	-2.887	44745.501	2	3d ⁵ (² F)4s	b ³ F	79458.450	2	3d ⁵ (a ² D)4p	x ³ F ^o	obs weak
2891.658	-3.178	66929.517	2	3d ⁵ (⁴ P)4p	z ⁵ S ^o	101501.621	3	3d ⁵ (⁴ G)5s	e ⁵ G	obs at 2891.65
2900.889	-3.778	62587.547	3	3d ⁵ (² D)4s	d ³ D	97049.640	4	3d ⁴ (⁵ D)4s4p(³ P ^o)	s ³ F ^o	obs
2918.568	-5.863	55697.014	4	3d ⁴ 4s ²	c ⁵ D	89950.370	3	3d ⁵ (a ² G)4p	x ¹ F ^o	obs
2931.882	-4.200	39813.760	2	3d ⁵ (⁴ D)4s	b ³ D	73911.570	1	3d ⁵ (⁴ P)4p	z ³ S ^o	pred weak
2932.315	-4.266	32836.740	1	3d ⁵ (⁴ D)4s	b ⁵ D	66929.517	2	3d ⁵ (⁴ P)4p	z ⁵ S ^o	STRONG, pred weak